
Introduction to the Finite Element method

Gustavo C. Buscaglia

ICMC-USP, São Carlos, Brasil
gustavo.buscaglia@gmail.com

Motivation

- For **elliptic** and **parabolic** problems, the most popular approximation method is the FEM.
- It is **general**, not restricted to linear problems, or to isotropic problems, or to any subclass of mathematical problems.
- It is **geometrically flexible**, complex domains are quite easily treated, not requiring adaptations of the method itself.
- It is **easy to code**, and the coding is quite problem-independent. Boundary conditions are much easier to deal with than in other methods.
- It is **robust**, because in most cases the mathematical problem has an underlying variational structure (energy minimization, for example).

Overview

- **Galerkin approximations:** Differential, variational and extremal formulations of a simple 1D boundary value problem. Well-posedness of variational formulations. Functional setting. Strong and weak coercivity. Lax-Milgram lemma. Banach's open mapping theorem. Céa's best-approximation property. Convergence under weak coercivity. (2 lectures)
- **The spaces of FEM and their implementation:** (3 lectures)
- **Interpolation error and convergence:** (2 lectures)
- **Application to convection-diffusion-reaction problems:** (2 lectures)
- **Application to linear elasticity:** (2 lectures)
- **Mixed problems:** (2 lectures)
- **FEM for parabolic problems:** (2 lectures)

1 Galerkin approximations

1.1 Variational formulation of a simple 1D example

Let u be the solution of

$$\begin{cases} -u'' + u = f & \text{in } (0, 1) \\ u(0) = u(1) = 0 \end{cases} \quad (1.1)$$

The **differential formulation** (DF) of the problem requires $-u'' + u$ to be exactly equal to f in **all** points $x \in (0, 1)$.

Multiplying the equation by any function v and integrating by parts (recall that

$$\int_0^1 w' z \, dx = w(1)z(1) - w(0)z(0) - \int_0^1 w z' \, dx \quad (1.2)$$

holds for all w and z that are *regular enough*) one obtains that u satisfies

$$\int_0^1 (u' v' + u v) \, dx - u'(1)v(1) + u'(0)v(0) = \int_0^1 f v \, dx \quad \forall v. \quad (1.3)$$

- The requirement “for all x ” of the DF has become “for all functions v ”.
- Does equation (1.3) fully determine u ?
- What happened with the boundary conditions?

Consider the following problem in **variational formulation** (VF): “Determine $u \in W$, such that $u(0) = u(1) = 0$ and that

$$\int_0^1 (u' v' + u v) dx = \int_0^1 f v dx \quad (1.4)$$

holds for all $v \in W$ satisfying $v(0) = v(1) = 0$.”

Prop. 1.1 *The solution u of the DF (eq. 1.1) is also a solution of the VF if W consists of continuous functions of sufficient regularity. As a consequence, problem VF admits at least one solution whenever DF does.*

Proof. Following the steps that lead to the VF, it becomes clear that the only requirement for u to satisfy (1.4) is that the integration by parts formula (1.2) be valid. \square

Exo. 1.1 *Show that the solution of*

$$\begin{cases} -u'' + u = f & \text{in } (0, 1) \\ u(0) = 0, \quad u'(1) = g \in \mathbb{R} \end{cases} \quad (1.5)$$

is a solution to: “Find $u \in W$ such that $u(0) = 0$ and that

$$\int_0^1 (u' v' + u v) dx = \int_0^1 f v dx + g v(1) \quad (1.6)$$

holds for all $v \in W$ satisfying $v(0) = 0$.”

Consider the following problem in **extremal formulation** (EF): “Determine $u \in W$ such that it minimizes the function

$$J(w) = \int_0^1 \left(\frac{1}{2}w'(x)^2 + \frac{1}{2}w(x)^2 - f w \right) dx \quad (1.7)$$

over the functions $w \in W$ that satisfy $w(0) = w(1) = 0$.”

Prop. 1.2 *The unique solution u of (1.1) is also a solution to EF. As a consequence, EF admits at least one solution.*

Proof. We need to show that $J(w) \geq J(u)$ for all $w \in W_0$, where

$$W_0 = \{w \in W, w(0) = w(1) = 0\}$$

Writing $w = u + \alpha v$ and replacing in (1.7) one obtains

$$J(u + \alpha v) = J(u) + \alpha \left[\int_0^1 (u' v' + u v - f v) dx \right] + \alpha^2 \int_0^1 \left(\frac{1}{2}v'(x)^2 + \frac{1}{2}v(x)^2 \right) dx$$

The last term is not negative and the second one is zero. \square

Exo. 1.2 *Identify the EF of the previous exercise.*

Prop. 1.3 Let u be the solution of

$$\begin{cases} -u'' + u = f & \text{in } (0, 1) \\ u(0) = 1, \quad u'(1) = g \in \mathbb{R} \end{cases} \quad (1.8)$$

then u is also a solution of “Determine $u \in W$ such that $u(0) = 1$ and that

$$\int_0^1 (u' v' + u v) dx = \int_0^1 f v dx + g v(1) \quad (1.9)$$

holds for all $v \in W$ satisfying $v(0) = 0$.”

Further, defining for any $a \in \mathbb{R}$

$$W_a = \{w \in W, w(0) = a\},$$

u minimizes over W_1 the function

$$J(w) = \int_0^1 \left(\frac{1}{2} w'(x)^2 + \frac{1}{2} w(x)^2 - f w \right) dx - g w(1). \quad (1.10)$$

Exo. 1.3 Prove the last proposition.

Let us define the bilinear and linear forms corresponding to problem (1.1):

$$a(v, w) = \int_0^1 (v'w' + vw) dx \qquad \ell(v) = \int_0^1 f v dx \qquad (1.11)$$

and the function $J(v) = \frac{1}{2}a(v, v) - \ell(v)$. Remember that W is a space of functions with some (yet unspecified) regularity and let $W_0 = \{w \in W, w(0) = w(1) = 0\}$.

The three formulations that we have presented up to now are, thus:

DF: Find a function u such that

$$-u''(x) + u(x) = f(x) \quad \forall x \in (0, 1), \qquad u(0) = u(1) = 0$$

VF: Find a function $u \in W_0$ such that

$$a(u, v) = \ell(v) \quad \forall v \in W_0$$

EF: Find a function $u \in W_0$ such that

$$J(u) \leq J(w) \quad \forall w \in W_0$$

and we know that the exact solution of DF is also a solution of VF and of EF.

The logic of the construction is justified by the following

Theorem 1.4 *If W is taken as*

$$W = \{w : (0, 1) \rightarrow \mathbb{R}, \int_0^1 w(x)^2 dx < +\infty, \int_0^1 w'(x)^2 dx < +\infty\} \stackrel{\text{def}}{=} H^1(0, 1)$$

and if f is such that there exists $C \in \mathbb{R}$ for which

$$\int_0^1 f(x) w(x) dx \leq C \sqrt{\int_0^1 w'(x)^2 dx} \quad \forall w \in W_0 \quad (1.12)$$

then problems (VF) and (EF) have one and only one solution, and their solutions coincide.

The proof will be given later, now let us consider its consequences:

- The differential equation has at most one solution in W .
- If the solution u to (VF)-(EF) is regular enough to be considered a solution to (DF), then u is the solution to (DF).
- If the solution u to (VF)-(EF) is not regular enough to be considered a solution to (DF), then (DF) has no solution.

\Rightarrow (VF) is a generalization of (DF).

Exo. 1.4 Show that $W_0 \subset C^0(0, 1)$. Further, compute $C \in \mathbb{R}$ such that

$$\max_{x \in [0, 1]} |w(x)| \leq C \sqrt{\int_0^1 w'(x)^2 dx} \quad \forall w \in W_0$$

Hint: You may assume that $\int_0^1 f(x)g(x) dx \leq \sqrt{\int_0^1 f(x)^2 dx} \sqrt{\int_0^1 g(x)^2 dx}$ for any f and g (Cauchy-Schwarz).

Exo. 1.5 Consider $f(x) = |x - 1/2|^\gamma$. For which exponents γ is $\int_0^1 f(x)w(x) dx < +\infty$ for all $w \in W_0$?

Exo. 1.6 Consider as f the “Dirac delta function” at $x = 1/2$, that we will denote by $\delta_{1/2}$. It can be considered as a “generalized” function defined by

$$\int_0^1 \delta_{1/2}(x) w(x) dx = w(1/2) \quad \forall w \in C^0(0, 1)$$

Prove that $\delta_{1/2}$ satisfies (1.12) and determine the analytical solution to (VF).

Exo. 1.7 Determine the DF and the EF corresponding to the following VF: “Find $u \in W = H^1(0, 1)$, $u(0) = 1$, such that

$$\int_0^1 (u'w' + uw) dx = w(1/2) \quad \forall w \in W_0 \tag{1.13}$$

where $W_0 = \{w \in W, w(0) = 0\}$.”

1.2 Variational formulations in general

Let V be a Hilbert space with norm $\|\cdot\|_V$. Let $a(\cdot, \cdot)$ and $\ell(\cdot)$ be bilinear and linear forms on V satisfying (continuity), for all $v, w \in V$,

$$a(v, w) \leq N_a \|v\|_V \|w\|_V, \quad \ell(v) \leq N_\ell \|v\|_V \quad (1.14)$$

This last inequality means that $\ell \in V'$, the (topological) dual of V . The minimum N_ℓ that satisfies this inequality is called the norm of ℓ in V' , i.e.

$$\|\ell\|_{V'} \stackrel{\text{def}}{=} \sup_{0 \neq v \in V} \frac{\ell(v)}{\|v\|_V} \quad (1.15)$$

The abstract VF we consider here is:

$$\text{“Find } u \in V \text{ such that } \quad a(u, v) = \ell(v) \quad \forall v \in V\text{”} \quad (1.16)$$

Exo. 1.8 Assume that V is finite dimensional, of dimension n , and let $\{\phi^1, \phi^2, \dots, \phi^n\}$ be a basis. Show that (1.16) is then equivalent to

$$\underline{V}^T \underline{A} \underline{U} = \underline{V}^T \underline{L} \quad \forall \underline{V} \in \mathbb{R}^n, \quad (1.17)$$

which in turn is equivalent to the linear system

$$\underline{A} \underline{U} = \underline{L}; \quad (1.18)$$

where

$$A_{ij} \stackrel{\text{def}}{=} a(\phi^j, \phi^i), \quad L_i \stackrel{\text{def}}{=} \ell(\phi^i) \quad (1.19)$$

and \underline{U} is the coefficient column vector of the expansion of u , i.e.,

$$u = \sum_{i=1}^n U_i \phi^i \quad (1.20)$$

Def. 1.5 The bilinear form $a(\cdot, \cdot)$ is said to be **strongly coercive** if there exists $\alpha > 0$ such that

$$a(v, v) \geq \alpha \|v\|_V^2 \quad \forall v \in V \quad (1.21)$$

Def. 1.6 The bilinear form $a(\cdot, \cdot)$ is said to be **weakly coercive** (or to satisfy an **inf-sup** condition) if there exists $\beta > 0$ such that

$$\sup_{0 \neq w \in V} \frac{a(v, w)}{\|w\|_V} \geq \beta \|v\|_V \quad \forall v \in V \quad (1.22)$$

and

$$\sup_{0 \neq v \in V} \frac{a(v, w)}{\|v\|_V} \geq \beta \|w\|_V \quad \forall w \in V \quad (1.23)$$

Exo. 1.9 Prove that strong coercivity implies weak coercivity.

Exo. 1.10 Prove that, if V is finite dimensional, then **(i)** $a(\cdot, \cdot)$ is strongly coercive iff $\underline{\underline{A}}$ is positive definite ($\underline{\underline{X}}^T \underline{\underline{A}} \underline{\underline{X}} > 0 \forall \underline{\underline{X}} \in \mathbb{R}^n$), and **(ii)** $a(\cdot, \cdot)$ is weakly coercive iff $\underline{\underline{A}}$ is invertible.

Exo. 1.11 Prove that, if $a(\cdot, \cdot)$ is weakly coercive, then the solution u of (1.16) depends continuously on the forcing $\ell(\cdot)$. Specifically, prove that

$$\|u\|_V \leq \frac{1}{\beta} \|\ell\|_{V'} \quad (1.24)$$

Theorem 1.7 *Assuming V to be a Hilbert space, problem (1.16) is well posed for any $\ell \in V'$ if and only if (i) $a(\cdot, \cdot)$ is continuous, and (ii) $a(\cdot, \cdot)$ is weakly coercive.*

A simpler version of this result is known as **Lax-Milgram lemma**:

Theorem 1.8 *Assuming V to be a Hilbert space, if $a(\cdot, \cdot)$ is continuous and strongly coercive then problem (1.16) is well posed for any $\ell \in V'$.*

Proof. This proof uses the so-called “Galerkin method”, which will be useful to introduce... the Galerkin method!

Let $\{\phi^i\}$ be a basis of V . Denoting $V_N = \text{span}(\phi^1, \dots, \phi^N)$ we can define $u_N \in V_N$ as the unique solution of $a(u_N, v) = \ell(v)$ for all $v \in V_N$. This generates a sequence $\{u_N\}_{N=1,2,\dots}$ in V . Further, this sequence is bounded, because

$$\|u_N\|_V^2 \leq \frac{1}{\alpha} a(u_N, u_N) = \frac{1}{\alpha} \ell(u_N) \leq \frac{\|\ell\|_{V'}}{\alpha} \|u_N\|_V \quad \Rightarrow \quad \|u_N\|_V \leq \frac{\|\ell\|_{V'}}{\alpha}, \quad \forall N$$

Recalling the weak compactness of bounded sets in Hilbert spaces, there exists $u \in V$ such that a subsequence of $\{u_N\}$ (still denoted by $\{u_N\}$ for simplicity) converges to u weakly. It remains to prove that $a(u, v) = \ell(v)$ for all $v \in V$. To see this, notice that

$$a(u, \phi^i) = a(\lim_N u_N, \phi^i) = \lim_N a(u_N, \phi^i) = \ell(\phi^i)$$

where the last equality holds because $a(u_N, \phi^i) = \ell(\phi^i)$ whenever $N \geq i$. Uniqueness is left as an exercise. \square

Exo. 1.12 *Prove uniqueness in the previous theorem (bounded sequences may have several accumulation points).*

Remark 1.9 The space $L^2(a, b)$ (also denoted by $H^0(a, b)$) is the Hilbert space of functions $f : (a, b) \rightarrow \mathbb{R}$ such that $\int_a^b f^2(x) dx < +\infty$.

The scalar product is

$$(f, g)_{L^2(a,b)} = \int_a^b f(x)g(x) dx \quad (1.25)$$

and accordingly

$$\|f\|_{L^2(a,b)} = (f, f)_{L^2(a,b)}^{1/2} = \sqrt{\int_a^b f^2(x) dx} . \quad (1.26)$$

Also of frequent use are the Hilbert spaces $H^1(a, b)$ and $H^2(a, b)$:

$$H^1(a, b) = \{f \in L^2(a, b) \mid f' \in L^2(a, b)\} \quad (1.27)$$

$$|f|_{H^1(a,b)} = \|f'\|_{L^2(a,b)} \quad (1.28)$$

$$\|f\|_{H^1(a,b)} = \|f\|_{L^2(a,b)} + |f|_{H^1(a,b)} \quad (1.29)$$

$$H^2(a, b) = \{f \in H^1(a, b) \mid f'' \in L^2(a, b)\} \quad (1.30)$$

$$|f|_{H^2(a,b)} = \|f''\|_{L^2(a,b)} \quad (1.31)$$

$$\|f\|_{H^2(a,b)} = \|f\|_{H^1(a,b)} + |f|_{H^2(a,b)} \quad (1.32)$$

Exo. 1.13 Other equivalent norms can be defined in $H^1(a, b)$, e.g.,

1. $\| \|f\| \|_{H^1(a,b)} = \left(\|f\|_{L^2(a,b)}^2 + |f|_{H^1(a,b)}^2 \right)^{1/2}$

2. $\| \|f\| \|_{H^1(a,b)} = \max \left(\|f\|_{L^2(a,b)}, |f|_{H^1(a,b)} \right)$

3. $\| \|f\| \|_{H^1(a,b)} = \|f\|_{L^2(a,b)} + \|\ell f'\|_{L^2(a,b)}$, where $\ell : (a, b) \rightarrow \mathbb{R}$ satisfies $0 < \ell_{\min} \leq \ell(x) \leq \ell_{\max}$ for all $x \in (a, b)$. Notice that if $\ell(x)$ has dimensions of length then this norm is unit-consistent.

Find the constants c and C such that $c\|f\| \leq \| |f| \| \leq C\|f\|$.

Remark 1.10 For the spaces $H^1(a, b)$ and $H^2(a, b)$ to be complete, one needs a weaker definition of the derivative. For this purpose, one first introduces the space

$$\mathcal{D}(a, b) = C_0^\infty(a, b) = \{ \varphi \in C^\infty(a, b) \mid \varphi \text{ has compact support in } (a, b) \} . \quad (1.33)$$

Given a function $f : (a, b) \rightarrow \mathbb{R}$, if there exists $g : (a, b) \rightarrow \mathbb{R}$ such that

$$\int_a^b g(x) \varphi(x) dx = - \int_a^b f(x) \varphi'(x) dx , \quad \forall \varphi \in \mathcal{D}(a, b) , \quad (1.34)$$

then we say that f' exists **in a weak sense**, and that $f' = g$.

Exo. 1.14 Show that the function

$$\phi(x) = \begin{cases} \exp(1/(|x|^2 - 1)) & \text{if } |x| < 1 \\ 0 & \text{if } |x| \geq 1 \end{cases} \quad (1.35)$$

belongs to $\mathcal{D}(\mathbb{R})$. By suitably shifting and scaling the argument of ϕ show that $\mathcal{D}(a, b)$ has infinite dimension for all $a < b$. (Hint: See Brenner-Scott, p. 27)

Exo. 1.15 Consider $f(x) = 1 - |x|$ in the domain $(-1, 1)$. Prove that its weak derivative is given by

$$f'(x) = \begin{cases} 1 & \text{if } x < 0 \\ -1 & \text{if } x > 0 \end{cases} . \quad (1.36)$$

Prove also that f'' does not exist. (Hint: See Brenner-Scott, p. 28)

Exo. 1.16 Let $f \in L^2(a, b)$, and let $V = H^1(a, b)$. Show that $\ell(v) = \int_a^b f(x) v(x) dx$ belongs to V' and that $\|\ell\|_{V'} \leq \|f\|_{L^2(a, b)}$.

1.3 Galerkin approximations

The previous proof suggests a numerical method, the Galerkin method, to approximate the solution of a variational problem and thus of an elliptic PDE. The idea is simply to restrict the variational problem to a subspace of V that we will denote by V_h .

Discrete variational problem (Galerkin): Find $u_h \in V_h$ such that

$$a(u_h, v_h) = \ell(v_h) \quad \forall v_h \in V_h \quad (1.37)$$

When the bilinear form $a(\cdot, \cdot)$ is symmetric and strongly coercive, this discrete problem is equivalent to

Discrete extremal problem (Galerkin): Find $u_h \in V_h$ which minimizes over V_h the function

$$J(w) = \frac{1}{2} a(w, w) - \ell(w) \quad (1.38)$$

Exo. 1.17 *Prove this last assertion.*

The natural questions that arise are:

- Does u_h exist? Is it unique?
- Does u_h approximate u (the exact solution)?
- How difficult is it to compute u_h ?

Does u_h exist? Is it unique?

Case 1) Strong coercivity of the form $a(\cdot, \cdot)$ over V

If $a(\cdot, \cdot)$ is strongly coercive over V , then

$$\inf_{0 \neq w \in V} \frac{a(w, w)}{\|w\|_V^2} = \alpha > 0.$$

If $V_h \subset V$, then $a(\cdot, \cdot)$ is strongly coercive over V_h (because the infimum is taken over a smaller set). Then u_h exists and is unique as a consequence of Exo. 1.10.

Case 2) Weak coercivity of the form $a(\cdot, \cdot)$ over V

If $a(\cdot, \cdot)$ is just weakly coercive over V , then it may or may not be weakly coercive over V_h . Compare the two following conditions

$$(A) \inf_{w \in V} \sup_{v \in V} \frac{a(w, v)}{\|w\|_V \|v\|_V} = \beta > 0, \quad (B) \inf_{w \in V_h} \sup_{v \in V_h} \frac{a(w, v)}{\|w\|_V \|v\|_V} = \beta_h > 0.$$

It is not true that $(A) \Rightarrow (B)$ because the sup in (B) is taken over a smaller set. In this case the weak coercivity of the discrete problem must be proven independently, it is not inherited from the weak coercivity over the whole space V .

Does u_h approximate u ?

Case 1) Strong coercivity of the form $a(\cdot, \cdot)$ over V

Lemma 1.11 (J. Céa) *If $a(\cdot, \cdot)$ and $\ell(\cdot)$ are continuous in V and $a(\cdot, \cdot)$ is strongly coercive, then*

$$\|u - u_h\|_V \leq \frac{N_a}{\alpha} \|u - v_h\|_V \quad \forall v_h \in V_h \quad (1.39)$$

Proof. Notice the so-called **Galerkin orthogonality**:

$$a(u - u_h, v_h) = 0 \quad \forall v_h \in V_h \quad (1.40)$$

which implies that $a(u - u_h, u - u_h) = a(u - u_h, u - v_h)$ for all $v_h \in V_h$. Using this,

$$\|u - u_h\|_V^2 \leq \frac{1}{\alpha} a(u - u_h, u - u_h) = \frac{1}{\alpha} a(u - u_h, u - v_h) \leq \frac{N_a}{\alpha} \|u - u_h\|_V \|u - v_h\|_V \quad \forall v_h \in V_h$$

In other words, $\|u - u_h\|_V \leq C \inf_{v_h \in V_h} \|u - v_h\|_V$. \square

Let h be a real parameter, typically a “mesh size”. We say that a family $\{V_h\}_{h>0} \subset V$ satisfies the **approximability property** if:

$$\lim_{h \rightarrow 0} \text{dist}(u, V_h) = \lim_{h \rightarrow 0} \inf_{v \in V_h} \|u - v\|_V = 0 \quad (1.41)$$

Corollary 1.12 *If $a(\cdot, \cdot)$ and $\ell(\cdot)$ are continuous in V , $a(\cdot, \cdot)$ is strongly coercive, and the family $\{V_h\}_{h>0} \subset V$ satisfies (1.41), then*

$$\lim_{h \rightarrow 0} u_h = u$$

in the sense of the norm $\|\cdot\|_V$.

If the strongly coercive bilinear form is **symmetric**, then $a(\cdot, \cdot)$ is a **scalar product** over V . In this case, Galerkin orthogonality corresponds to: **The Galerkin solution u_h is the orthogonal projection of u onto V_h .**

Further, the **energy norm** can be defined

$$\|v\|_a = \sqrt{a(v, v)}, \quad (1.42)$$

which satisfies the equivalence

$$\alpha^{\frac{1}{2}} \|v\|_V \leq \|v\|_a \leq N_a^{\frac{1}{2}} \|v\|_V. \quad (1.43)$$

Exo. 1.18 Show that the Galerkin approximation is **optimal** in the energy norm,

$$\|u - u_h\|_a \leq \|u - v_h\|_a, \quad \forall v_h \in V_h, \quad (1.44)$$

without the constants that appear in Céa's lemma. Further show that the following sharper estimate holds:

$$\|u - u_h\|_V \leq \left(\frac{N_a}{\alpha}\right)^{\frac{1}{2}} \|u - v_h\|_V, \quad \forall v_h \in V_h. \quad (1.45)$$

Case 2) Weak coercivity of the form $a(\cdot, \cdot)$ over V_h

Assume now that the weak coercivity constant β_h is positive for all $h > 0$, so that u_h exists and is unique. Notice that Galerkin orthogonality still holds.

Lemma 1.13 *If $a(\cdot, \cdot)$ and $\ell(\cdot)$ are continuous in V , and $a(\cdot, \cdot)$ is weakly coercive in V_h with constant $\beta_h > 0$, then*

$$\|u - u_h\|_V \leq \left(1 + \frac{N_a}{\beta_h}\right) \|u - v_h\|_V \quad \forall v_h \in V_h \quad (1.46)$$

Proof. One begins by decomposing the error as follows (we omit the subindex V in the norm)

$$\|u - u_h\| \leq \|u - v_h\| + \|u_h - v_h\| \quad \forall v_h \in V_h \quad (1.47)$$

and then using the weak coercivity

$$\|u_h - v_h\| \leq \frac{1}{\beta_h} \sup_{w_h \in V_h} \frac{a(u_h - v_h, w_h)}{\|w_h\|} = \frac{1}{\beta_h} \sup_{w_h \in V_h} \frac{a(u - v_h, w_h)}{\|w_h\|} \leq \frac{N_a}{\beta_h} \|u - v_h\|$$

Substituting this into (1.47) one proves the claim. \square

Corollary 1.14 *Under the hypotheses of Lemma 1.13, if there exists $\beta_0 > 0$ such that $\beta_h > \beta_0$ for all h and the family $\{V_h\}_{h>0} \subset V$ satisfies (1.41), then*

$$\lim_{h \rightarrow 0} u_h = u$$

in the sense of the norm $\|\cdot\|_V$.

How difficult is it to compute u_h ?

Let us go back to our problem $-u'' + u = f$ in $(0, 1)$ with $u(0) = u(1) = 0$, which in VF requires to compute $u \in H^1(0, 1)$ satisfying the boundary conditions and such that

$$\int_0^1 [u'(x)v'(x) + u(x)v(x)] dx = \int_0^1 f(x)v(x) dx \quad (1.48)$$

Suitable spaces for the Galerkin approximation are, for example,

- \mathcal{P}_k : The polynomials of degree up to k .
- \mathcal{F}_k : The space generated by the functions $\phi^m(x) = \sin(m\pi x)$, $m = 1, 2, \dots, k$.

Exo. 1.19 Show that $a(\cdot, \cdot)$ is continuous and strongly coercive over $V = H^1(0, 1)$ with the norm

$$\|w\|_V \stackrel{\text{def}}{=} \left[\int_0^1 [w'(x)^2 + w(x)^2] dx \right]^{\frac{1}{2}}$$

Exo. 1.20 Build a small program in Matlab or Octave (or something else) that solves the Galerkin approximation of problem (1.48) considering $f = \delta_{1/4}$ and the spaces \mathcal{P}_k and/or \mathcal{F}_k , for some values of k . Compare the results to the analytical solution building plots of u and u_h . Also, build graphs of $\|u - u_h\|$ vs k .

In general, however, the construction of spaces of global basis functions, as the ones above, is not practical because it leads to dense matrices. In the next chapter we will introduce the spaces of the FEM, which are characterized by having bases with small support and thus lead to sparse matrices.

Exercises

Reading assignment: Read Chapter 1 of Duran's notes (all of it).

Exo. 1.21 Carry out the “easy computation” that shows that $\underline{\underline{A}}$ is the tridiagonal matrix such that the diagonal elements are $2/h + 2h/3$ and the extra-diagonal elements are $-1/h + h/6$ (Durán, page 3).

Exo. 1.22 Can a symmetric bilinear form be weakly coercive but not strongly coercive?

Exo. 1.23 To what variational formulation and what differential formulation corresponds the following extremal formulation?

Find $u \in V$, V consisting of functions that are smooth in $(0, 1/2)$ and $(1/2, 1)$ but can exhibit a (bounded) discontinuity at $x = 1/2$, that minimizes the function

$$J(w) = \int_0^1 [w'(x)^2 + 2w(x)^2] dx + 4 [w(1/2+) - w(1/2-)]^2 - \int_0^{1/2} 7 w(x) dx - 9w(0) \quad (1.49)$$

where $w(1/2\pm)$ represent the values on each side of the discontinuity. Notice that the space V (is it a vector space really?) has no boundary condition imposed. What are the boundary conditions of the DF at $x = 0$ and $x = 1$?

Exo. 1.24 Consider the bilinear form

$$a(u, v) = \int_0^1 u'(x) v'(x) dx.$$

Prove that this form is not strongly coercive in $H^1(0, 1)$ considering the norm

$$\|w\|_{H^1} \stackrel{\text{def}}{=} \left\{ \int_0^1 [u'(x)^2 + u(x)^2] dx \right\}^{\frac{1}{2}}$$

and that it is, with the same norm, in

$$H_0^1(0, 1) \stackrel{\text{def}}{=} \{w \in H^1(0, 1), w(0) = w(1) = 0\}.$$

1.4 Variational formulations in 2D and 3D

The ideas are similar, but we need another integration by parts formula:

Lemma 1.15 *Let $f : \Omega \rightarrow \mathbb{R}$ be an integrable function, with Ω a Lipschitz bounded open set in \mathbb{R}^d and $\partial_i f$ integrable over Ω , then*

$$\int_{\Omega} \partial_i f \, d\Omega = \int_{\partial\Omega} f n_i \, d\Gamma \quad (1.50)$$

Notice that this implies that

$$\int_{\Omega} \nabla \cdot \mathbf{v} \, d\Omega = \int_{\partial\Omega} \mathbf{v} \cdot \check{\mathbf{n}} \, d\Gamma \quad (1.51)$$

and that

$$\int_{\Omega} v \nabla^2 u \, d\Omega = \int_{\partial\Omega} v \nabla u \cdot \check{\mathbf{n}} \, d\Gamma - \int_{\Omega} \nabla v \cdot \nabla u \, d\Omega \quad (1.52)$$

We will also introduce the notation

Def. 1.16 *The Lebesgue space $L^p(\Omega)$, where $p \geq 1$, is the set of all functions such that their $L^p(\Omega)$ -norm is finite,*

$$\|w\|_{L^p(\Omega)} \stackrel{\text{def}}{=} \left[\int_{\Omega} |w(x)|^p \, dx \right]^{\frac{1}{p}} \quad (1.53)$$

Exa. 1.17 (Poisson equation) Consider the DF

$$-\nabla^2 u = f \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega \quad (1.54)$$

where ∇ is the gradient operator and $\nabla^2 u = \sum_{i=1}^d \partial_{ii}^2 u$.

A suitable variational formulation is: Find $u \in V$ such that

$$a(u, v) = \ell(v) \quad \forall v \in V$$

where

$$a(u, v) = \int_{\Omega} \nabla u \cdot \nabla v \, d\Omega, \quad \ell(v) = \int_{\Omega} f v \, d\Omega \quad \text{and} \quad (1.55)$$

$$V = H_0^1(\Omega) = \{w \in L^2(\Omega), \partial_i w \in L^2(\Omega) \forall i = 1, \dots, d, w = 0 \text{ on } \partial\Omega\}$$

which is a Hilbert space with the norm

$$\|w\|_{H^1} = (\|w\|_{L^2}^2 + \|\nabla w\|_{L^2}^2)^{\frac{1}{2}} \quad (1.56)$$

Exo. 1.25 Prove that if u is a solution of the DF, then it solves the VF.

Exo. 1.26 Prove that $a(\cdot, \cdot)$ is continuous in V . Prove that $\ell(\cdot)$ is continuous in V if $f \in L^2(\Omega)$. Is this last condition necessary?

Exo. 1.27 Determine the EF of the Poisson problem.

Exo. 1.28 Is $a(\cdot, \cdot)$ strongly coercive?

Exo. 1.29 Let Ω be the unit circle. Determine for which exponents γ is the function r^γ in $H^1(\Omega)$.

Exo. 1.30 Assume that the domain Ω is divided into subdomains Ω_1 and Ω_2 by a smooth internal boundary Γ . Let V consist of functions such that their restrictions to Ω_i belong to $H^1(\Omega_i)$ and that are continuous across Γ . Determine the VF corresponding to the following EF: Find $u \in V$ that minimizes

$$J(w) = \int_{\Omega_1} \frac{w^2 + \|\nabla w\|^2}{2} d\Omega + \int_{\Omega_2} \frac{3\|\nabla w\|^2}{2} d\Omega + \int_{\Gamma} (5w^2 - w) d\Gamma$$

over V .

Exo. 1.31 Determine the DF that corresponds to the previous exercise.

2 Finite element spaces and interpolation

The basic reference for what follows is Ciarlet [?]. Basically, the idea is to define finite element spaces that are locally polynomial and that contain complete polynomials of degree k in the space variables. With a judicious choice of the nodes (degrees of freedom), these piecewise polynomial functions can be made continuous by construction (if needed).

In the previous chapter it was shown that if there exists $\beta > 0$ such that, for all $w_h \in V_h$ and all $h > 0$,

$$\sup_{v_h \in V_h} \frac{a(w_h, v_h)}{\|v_h\|_V} \geq \beta \|w_h\|_V \quad (2.1)$$

then there exists $C > 0$ such that

$$\|u - u_h\|_V \leq C \inf_{v_h \in V_h} \|u - v_h\|_V \quad (2.2)$$

Notice that (2.1) is automatically satisfied if the bilinear form $a(\cdot, \cdot)$ is strongly coercive.

Denoting by $\mathcal{I}_h u$ the element-wise Lagrange interpolant of $u \in V \cap C^0(\overline{\Omega})$, it is obvious from (2.2) that

$$\|u - u_h\|_V \leq C \|u - \mathcal{I}_h u\|_V \quad (2.3)$$

The goal of this section is to introduce estimates of the interpolation error $\|u - \mathcal{I}_h u\|_V$ for some spaces V that appear in the applications.

2.1 Basic definitions

Def. 2.1 *A finite element in \mathbb{R}^n is a triplet (K, P_K, Σ_K) where*

- (i) *K is a closed (bounded) subset of \mathbb{R}^n with a nonempty interior and Lipschitz boundary;*
- (ii) *P_K is a finite-dimensional space of functions defined in K , of dimension m ;*

(iii) Σ_K is a set of m linear forms $\{\sigma_i\}_{i=1,\dots,m}$ which is P_K -unisolvent; i.e., if $p \in P_K$ then

$$\sigma(p) = 0 \quad \forall \sigma \in \Sigma_K \quad \Rightarrow \quad p = 0$$

It is implicitly assumed that the finite element is viewed with a larger function space $V(K)$ associated to it, in general a Sobolev space. Each $\sigma_i \in \Sigma_K$ is then assumed to be extended as an element of $V(K)'$.

Prop. 2.2 *There exists a basis $\{\mathcal{N}_i\}$ such that $\sigma_i(\mathcal{N}_j) = \delta_{ij}$.*

Whenever needed, we will write $\sigma_{K,i}$ instead of σ_i and $\mathcal{N}_{K,i}$ instead of \mathcal{N}_i to make explicit the element K being considered.

Def. 2.3 *If the degrees of freedom correspond to nodal values of the functions in $V(K)$ the element is called a Lagrange finite element. In this case, there exist X^1, \dots, X^m in K such that $\sigma_i(v) = v(X^i)$ for all $i = 1, \dots, m$.*

Exa. 2.4 *P_k elements.*

Finite elements are usually built by mapping a unique *master element* $(\widehat{K}, \widehat{P}, \widehat{\Sigma})$ onto (K, P_K, Σ_K) in a clever way. We denote by $\{\widehat{\sigma}_i\}$ the degrees of freedom of the master, and $\{\widehat{\mathcal{N}}_i\}$ the corresponding basis functions.

One begins by defining a linear bijective transformation

$$T_K : V(K) \rightarrow V(\widehat{K}) , \tag{2.4}$$

mapping functions defined on K onto functions defined on \widehat{K} . Its inverse, T_K^{-1} allows one to build P_K , i.e.,

$$P_K = \{T_K^{-1}\widehat{p}, \widehat{p} \in \widehat{P}\} . \tag{2.5}$$

Exo. 2.1 Two elements \widehat{K} , K , are said to be affine equivalent if there exists a bijective mapping $F_K : \widehat{K} \rightarrow K$ of the form

$$F_K(\hat{x}) = A_K \hat{x} + b . \quad (2.6)$$

Show that if $\widehat{P} = \mathbb{P}_k$ and T_K is defined by

$$(T_K v)(\hat{x}) = v(F_K(\hat{x})) \quad (2.7)$$

then $P_K = \mathbb{P}_k$.

This preservation of polynomial spaces makes the analysis of affine-equivalent elements much easier, but if F_K is not affine one still uses (2.7) for the definition of P_K in Lagrange finite elements (P_K will not consist of polynomials).

Prop. 2.5 If $(\widehat{K}, \widehat{P}, \widehat{\Sigma})$ is a master finite element and $T_K : V(K) \rightarrow V(\widehat{K})$ is linear and bijective, then the triplet (K, P_K, Σ_K) given by

$$K = F_K(\widehat{K}) \quad (2.8)$$

$$P_K = T_K^{-1} \widehat{P} \quad (2.9)$$

$$\Sigma_K = \{ \sigma_i \mid \sigma_i(p) = \alpha_i \widehat{\sigma}_i(T_K p), \forall p \in P_K \} \quad (2.10)$$

(where all α_i are non-zero) is a finite element. Further, the basis functions on K are given by

$$\mathcal{N}_i = \frac{1}{\alpha_i} T_K^{-1} \widehat{\mathcal{N}}_i . \quad (2.11)$$

In the case of Lagrange finite elements one takes $\alpha_i = 1$ and obtains

$$X^i = F_K(\widehat{X}^i) , \quad \sigma_i(p) = p(X^i) = p(F_K(\widehat{X}^i)) , \quad \mathcal{N}_i(F_K(\hat{x})) = \widehat{\mathcal{N}}_i(\hat{x}) . \quad (2.12)$$

Exo. 2.2 *Prove the previous proposition.*

Hint: One has to assume that \widehat{K} has Lipschitz boundary and that F_K is regular enough for $F_K(\widehat{K})$ to have Lipschitz boundary too. Because T_K is linear bijective, P_K will be a vector space of the same dimension as \widehat{P} . It remains to show that Σ_K is unisolvent. Let $p \in P_K$ such that $\sigma_i(p) = 0$ for all $i = 1, \dots, m$. Then $\widehat{\sigma}_i(T_K p) = 0$ for all i and thus $T_K p = 0$ because $\widehat{\Sigma}$ is unisolvent. The last assertion follows from $\sigma_i(\mathcal{N}_j) = \delta_{ij}$ and the case of Lagrange finite elements is a particular case of the former.

A case in which the scaling factors α_i in the previous proposition are needed is that of **Hermite finite elements**.

Exo. 2.3 *Build a basis for a cubic 1D Hermite finite element. For this, let K be an interval $[a, b]$, let $V(K) = H^2(K)$, $P_K = \mathbb{P}_3$ (cubic polynomials), and*

$$\Sigma_K = \{\theta_a, \theta_b, \eta_a, \eta_b\} , \quad (2.13)$$

where $\theta_a(v) = v(a)$, $\theta_b(v) = v(b)$, $\eta_a(v) = v'(a)$ and $\eta_b(v) = v'(b)$. Write down the basis functions.

Now consider the master element $\widehat{K} = [-1, 1]$ and the affine mapping

$$F_K(\hat{x}) = a + \frac{b-a}{2}(x+1) . \quad (2.14)$$

Defining T_K as in (2.7), find the factors $\{\alpha_i\}$ so that σ_i relates $\widehat{\sigma}_i$ according to (2.10). Write down the basis functions \widehat{N}_i and verify (2.11).

Raviart-Thomas finite element: The interpolation of vector fields in K with Lagrange finite elements is usually done one component at a time with the tools developed for scalar functions. A notable exception is the Raviart-Thomas element, very popular to approximate velocity fields in porous media. In this case $V(K)$ and P_K consist of **vector fields** and cannot be interpolated one component at a time.

Exo. 2.4 Let K be a simplex (triangle in 2D, tetrahedron in 3D), and let the space P_K be defined as

$$P_K = \mathbb{RT}_0 = (\mathbb{P}_0)^d \oplus x\mathbb{P}_0 , \quad (2.15)$$

which is of dimension $d + 1$. Defining as degrees of freedom the fluxes across each face (edge in 2D),

$$\sigma_i(p) = \int_{F_i} p \cdot \tilde{n} , \quad (2.16)$$

prove that $(K, P_K, \Sigma_K = \{\sigma_i\})$ is a finite element and that

$$\mathcal{N}_i(x) = \frac{1}{d \operatorname{meas}(K)} (x - a_i) , \quad (2.17)$$

where a_i is the vertex opposite to F_i .

The space V_K is in this case $H(\operatorname{div}, K)$ of vector fields in $L^2(K)^d$ with divergence in $L^2(K)$.

To obtain the RT0 element from a master element one needs a transformation $T_K : V(K) \rightarrow V(\hat{K})$ such that $P_K = T_K^{-1}\hat{P}$ and $\sigma_i(p) = \alpha_i \hat{\sigma}_i(T_K p)$ for all $p \in P_K$.

Let

$$\hat{\sigma}_i(\hat{p}) = \int_{\hat{F}_i} \hat{p} \cdot \hat{n} \, d\hat{F} , \quad (2.18)$$

and define the **Piola transformation**

$$\hat{v}(\hat{x}) = T_K v(\hat{x}) = \det(A_K) A_K^{-1} v(F_K(\hat{x})) . \quad (2.19)$$

Then

$$\hat{\sigma}_i(\hat{p}) = \det(A_K) \int_{\hat{F}_i} (A_K^{-1} p(F_K(\hat{x}))) \cdot \hat{n} \, d\hat{F} = \int_{\hat{F}_i} p(F_K(\hat{x})) \cdot (\det(A_K) A_K^{-T} \hat{n} \, d\hat{F}) \quad (2.20)$$

It is well-known that $\hat{n} \, d\hat{F}$ transforms as

$$\check{n} \, dF = \det(A_K) A_K^{-T} \hat{n} \, d\hat{F} , \quad (2.21)$$

so that, changing the integration variable to $x = F_K(\hat{x})$,

$$\hat{\sigma}_i(\hat{p}) = \int_{F_i} p(x) \cdot \check{n} \, dF = \sigma_i(p) . \quad (2.22)$$

The RT0 element is thus obtained from the master element using the Piola transformation as T_K and $\alpha_i = 1$.

Def. 2.6 *The local interpolation operator $\mathcal{I}_K : V(K) \rightarrow P_K$ is defined as*

$$\mathcal{I}_K v = \sum_{i=1}^m \sigma_i(v) \mathcal{N}_i \quad \forall v \in V(K)$$

Exo. 2.5 *This interpolation is indeed a projection:*

$$\mathcal{I}_K p = p \quad \text{for all } p \in P_K . \quad (2.23)$$

Exo. 2.6 *It is also preserved by composition with the T_K mapping:*

$$\widehat{\mathcal{I}_K v} = T_K \mathcal{I}_K v = \mathcal{I}_{\widehat{K}} T_K v = \mathcal{I}_{\widehat{K}} \widehat{v}, \quad \text{for all } v \in V(K) . \quad (2.24)$$

We now turn to the problem of estimating the interpolation error, i.e., $v - \mathcal{I}_K v$.

2.2 Local $L^\infty(K)$ estimates for P_1 -triangles

We begin by considering the case of P_1 -simplices (triangles in 2D, tetrahedra in 3D). It is a good exercise in which the estimates can be derived explicitly. It is also a good excuse to introduce the multi-point Taylor formula.

Theorem 2.7 *Let K be a P_1 -element, h_K its diameter and ρ_K the radius of the largest ball contained in K . Then, for all $v \in C^2(K)$,*

$$(a) \quad \|v - \mathcal{I}_K v\|_{L^\infty(K)} \leq \frac{d^2 h_K^2}{2} \max_{|\alpha|=2} \|D^\alpha v\|_{L^\infty(K)}$$

$$(b) \quad \max_{|\alpha|=1} \|D^\alpha(v - \mathcal{I}_K v)\|_{L^\infty(K)} \leq \frac{(d+1)d^2 h_K^2}{2\rho_K} \max_{|\alpha|=2} \|D^\alpha v\|_{L^\infty(K)}$$

Proof. Let X^j be the position of the j -th node of the element, then

$$\mathcal{I}_K v(x) = \sum_{j=1}^{d+1} v(X^j) \mathcal{N}^j(x) \quad (2.25)$$

We now perform a Taylor expansion *around* x , and evaluate it at X^j , obtaining

$$v(X^j) = v(x) + \sum_{k=1}^d \frac{\partial v}{\partial x_k}(x) (X_k^j - x_k) + \frac{1}{2} \sum_{k,\ell=1}^d \frac{\partial^2 v}{\partial x_k \partial x_\ell}(\xi) (X_k^j - x_k) (X_\ell^j - x_\ell) \quad (2.26)$$

where $\xi = \eta X^j + (1 - \eta)x$ for some $\eta \in [0, 1]$. Let us denote by $p^j(x)$ the second term in the right-hand side of (2.26), and by $r^j(x)$ the third term. By direct inspection we notice that

$$|r^j(x)| \leq \frac{d^2 h_K^2}{2} \max_{|\alpha|=2} \|D^\alpha v\|_{L^\infty(K)}$$

Let us now insert $v(X^j)$ from (2.26) into (2.25) to get

$$\mathcal{I}_K v(x) = \sum_{j=1}^{d+1} v(x) \mathcal{N}^j(x) + \sum_{j=1}^{d+1} p^j(x) \mathcal{N}^j(x) + \sum_{j=1}^{d+1} r^j(x) \mathcal{N}^j(x)$$

The first term on the right is equal to $v(x)$ because $\sum_j \mathcal{N}^j = 1$. The second term vanishes, since

$$\begin{aligned} \sum_{j=1}^{d+1} \sum_{k=1}^d \frac{\partial v}{\partial x_k}(x) (X_k^j - x_k) \mathcal{N}^j(x) &= \sum_{k=1}^d \frac{\partial v}{\partial x_k}(x) \left\{ \sum_{j=1}^{d+1} X_k^j \mathcal{N}^j(x) - x_k \sum_{j=1}^{d+1} \mathcal{N}^j(x) \right\} = \\ &= \sum_{k=1}^d \frac{\partial v}{\partial x_k}(x) \{x_k - x_k\} = 0 \end{aligned}$$

As a consequence, $v(x) - \mathcal{I}_K v(x) = \sum_{j=1}^{d+1} r^j(x) \mathcal{N}^j(x)$ and thus

$$|v(x) - \mathcal{I}_K v(x)| \leq \max_j |r^j(x)| \sum_j \mathcal{N}^j(x) = \max_j |r^j(x)| \leq \frac{d^2 h_K^2}{2} \max_{|\alpha|=2} \|D^\alpha v\|_{L^\infty(K)}$$

implying assertion (a). Now, by differentiating (2.25) and using (2.26) as before, one obtains

$$\frac{\partial \mathcal{I}_K v}{\partial x_m}(x) = \sum_j v(x) \frac{\partial \mathcal{N}^j}{\partial x_m}(x) + \sum_{j,k} \frac{\partial v}{\partial x_k}(x) (X_k^j - x_k) \frac{\partial \mathcal{N}^j}{\partial x_m}(x) + \sum_{j,k} r^j(x) \frac{\partial \mathcal{N}^j}{\partial x_m}(x)$$

On the right-hand side above, the first term vanishes and the second term happens to be equal to $\frac{\partial v}{\partial x_m}(x)$, since

$$\sum_{j,k} \frac{\partial v}{\partial x_k}(x) (X_k^j - x_k) \frac{\partial \mathcal{N}^j}{\partial x_m}(x) = \sum_k \frac{\partial v}{\partial x_k}(x) \left[\sum_j X_k^j \frac{\partial \mathcal{N}^j}{\partial x_m}(x) - x_m \sum_j \frac{\partial \mathcal{N}^j}{\partial x_m}(x) \right] =$$

$$= \sum_k \frac{\partial v}{\partial x_k}(x) \frac{\partial}{\partial x_m} \sum_j X_k^j \mathcal{N}^j(x) = \sum_k \frac{\partial v}{\partial x_k}(x) \frac{\partial x_k}{\partial x_m} = \frac{\partial v}{\partial x_m}(x)$$

As a consequence

$$\left| \frac{\partial \mathcal{I}_K v}{\partial x_m}(x) - \frac{\partial v}{\partial x_m}(x) \right| = \left| \sum_{j=1}^{d+1} r^j(x) \frac{\partial \mathcal{N}^j}{\partial x_m}(x) \right| \leq \max_j |r^j(x)| \sum_{j=1}^{d+1} \left| \frac{\partial \mathcal{N}^j}{\partial x_m}(x) \right|$$

The reader can convince himself that the norm of the gradient of a P_1 basis function, which equals one at one node and zero on the opposite side/face, can never be greater than $\frac{1}{\rho_K}$, which immediately leads to assertion (b). \square

2.3 Local estimates in Sobolev norms

The previous paragraph provides us with an interpolation estimate in the norm $L^\infty(K)$ for the function and its first derivatives. Most formulations studied so far, however, have $V = H^1(\Omega)$ and we need thus estimates of $u - \mathcal{I}_K u$ in the $H^m(K)$ -norm.

2.3.1 First estimates

A simplistic approach to estimate $\|u - \mathcal{I}_K u\|_{L^2(K)}$ for P_1 elements could be

$$\|u - \mathcal{I}_K u\|_{L^2(K)}^2 = \int_K (u - \mathcal{I}_K u)^2 \leq |K| \|u - \mathcal{I}_K u\|_{L^\infty(K)}^2 \leq 4|K| h_K^4 \max_{|\alpha|=2} \|D^\alpha u\|_{L^\infty(K)}^2$$

so that, with simplified notation,

$$\|u - \mathcal{I}_K u\|_{L^2(K)} \leq 2 \sqrt{|K|} h_K^2 \|D^2 u\|_{L^\infty(K)} \quad (2.27)$$

Proceeding analogously, we obtain a first estimate for $\|\nabla u - \nabla(\mathcal{I}_K u)\|_{L^2(K)}$,

$$\|\nabla u - \nabla(\mathcal{I}_K u)\|_{L^2(K)}^2 = \int_K \sum_{i=1}^d \left[\frac{\partial(u - \mathcal{I}_K u)}{\partial x_i} \right]^2 \leq |K| \sum_{i=1}^d \left\| \frac{\partial(u - \mathcal{I}_K u)}{\partial x_i} \right\|_{L^\infty(K)}^2$$

which from Th. 2.7 implies

$$\|\nabla u - \nabla(\mathcal{I}_K u)\|_{L^2(K)} \leq \sqrt{|K|} \frac{6 d h_K^2}{\rho_K} \|D^2 u\|_{L^\infty(K)} \quad (2.28)$$

Notice that these estimates require $u \in W^{2,\infty}(K)$, which is “too much” regularity.

Exo. 2.7 Consider the function $u(x) = |x|$ and its P_1 interpolant in the 1D simplex $K = (-h/2, h/2)$. Compute $\|u - \mathcal{I}_K u\|_{L^2(K)}$ and $\|u' - (\mathcal{I}_K u)'\|_{L^2(K)}$, compare to the previous estimates, and discuss briefly.

2.3.2 An L^2 -estimate without second derivatives

If the function to be interpolated does not have second derivatives in K , then $\|u - \mathcal{I}_K u\|_{L^2(K)}$ cannot be expected to be of order $\mathcal{O}(\sqrt{|K|} h_K^2)$. The following estimate, proved in *Buscaglia & Agouzal* (IMA J. Numer. Anal. 32, 672-686, 2012), has minimal requirements on both P_K and u . Notice in particular that P_K must contain the constants but not necessarily polynomials of degree 1.

Theorem 2.8 *Assume that the basis functions $\{\mathcal{N}^j\}$ ($j = 1, \dots, d+1$) of an element K satisfy: (H1) $\mathcal{N}^j(X^k) = \delta_{jk}$, (H2) $\sum_j \mathcal{N}^j(x) = 1$, (H3) $0 \leq \mathcal{N}^j(x) \leq 1$ for all j and for all $x \in K$.*

Then, for all $u \in W^{1,p}(K)$ with $p > d \geq 2$,

$$\|u - \mathcal{I}_K u\|_{L^2(K)} \leq \frac{p(d+1)}{p-d} |K|^{\frac{1}{2} - \frac{1}{p}} h_K \|\nabla u\|_{L^p(K)} \quad (2.29)$$

If ∇u is bounded we can take $p = +\infty$ to get

$$\|u - \mathcal{I}_K u\|_{L^2(K)} \leq (d+1) \sqrt{|K|} h_K \|\nabla u\|_{L^\infty(K)} \quad (2.30)$$

which is of order $\mathcal{O}(\sqrt{|K|} h_K)$.

2.3.3 Local interpolation estimates for Lagrange finite elements

Lagrange interpolation implies that the function being interpolated is at least in $C^0(K)$, since otherwise its nodal values would not be well defined.

Sobolev's imbedding theorems state that, for bounded convex domain K , $W^{m,p}(K) \subset C^0(K)$ if $mp > d$. Taking $p = 2$ (Hilbert spaces), m needs to be at least 1 in 1D and at least 2 in 2D/3D for $H^m(K)$ to consist of continuous functions.

Theorem 2.9 *Let (K, P_K, Σ_K) be a Lagrange finite element such that (a) P_K contains all polynomials of degree $\leq k$, and (b) it is affine-equivalent to the "master element" $(\widehat{K}, \widehat{P}, \widehat{\Sigma})$. Then, the Lagrange interpolant $\mathcal{I}_K u(x) = \sum_j u(X^j) \mathcal{N}^j(x)$ satisfies*

$$\|u - \mathcal{I}_K u\|_{L^2(K)} \leq C h_K^{\ell+1} \|D^{\ell+1} u\|_{L^2(K)} \quad (2.31)$$

for all $\ell \leq k$, with C depending on ℓ but not on h_K or u .

Similarly,

$$\|u - \mathcal{I}_K u\|_{H^1(K)} = \|\nabla u - \nabla(\mathcal{I}_K u)\|_{L^2(K)} \leq C \frac{h_K^{\ell+1}}{\rho_K} \|D^{\ell+1} u\|_{L^2(K)} \quad (2.32)$$

The proof of this theorem is somewhat involved. The interested reader may refer to Ciarlet [?] or to Ern-Guermond [?].

2.4 Global interpolation error

The obtention of global interpolation estimates is quite straightforward, but needs a few definitions.

2.4.1 Considerations about meshes

A mesh \mathcal{T}_h of a domain Ω in \mathbb{R}^d is a collection of compacts (elements) K_i , $i = 1, \dots, N_e$, such that

$$\bar{\Omega} = \bigcup_{i=1}^{N_e} K_i, \quad K_i \cap K_j = \emptyset \text{ if } i \neq j, \quad \partial\Omega \subset \bigcup_{i=1}^{N_e} \partial K_i \quad (2.33)$$

Def. 2.10 *The global interpolation operator $\mathcal{I}_h : W \rightarrow W_h$, where*

$$W = \{w \in L^1(\Omega), w|_K \in V(K), \forall K \in \mathcal{T}_h\}$$

$$W_h = \{w \in L^1(\Omega), w|_K \in P_K, \forall K \in \mathcal{T}_h\}$$

by

$$\mathcal{I}_h v = \sum_{K \in \mathcal{T}_h} \sum_i \sigma_{K,i}(v|_K) \mathcal{N}_{K,i} \quad (2.34)$$

Notice that, depending on the definition of the degrees of freedom, $\mathcal{I}_h v$ may be multiple-valued at element boundaries. The mesh is said to be conforming when $\mathcal{I}_h v$ belongs to the Sobolev space W in which the variational problem is posed.

The subscript h refers to the mesh size. In fact, in error estimates one has to consider not a single mesh but a family of meshes indexed by h , and study the error as $h \rightarrow 0$. The geometrical properties of the mesh refinement enter thus into consideration. Generally, the mesh-size parameter h is defined as

$$h = \max_{K \in \mathcal{T}_h} h_K \quad (2.35)$$

For global estimates in $H^m(\Omega)$ with $m \geq 1$ the ratio $s_K = \frac{h_K}{\rho_K}$ will appear. This motivates the definition of shape-regular (or, simply, regular) meshes:

Def. 2.11 *A family of meshes \mathcal{T}_h , parameterized by the parameter $h \in H$ (where H is some subset of \mathbb{R}), is said to be **shape-regular** if there exists $S \in \mathbb{R}$ such that*

$$s_K = \frac{h_K}{\rho_K} \leq S \quad \forall K \in \mathcal{T}_h, \quad \forall h \in H \quad (2.36)$$

A shape-regular mesh (rigorously speaking, family of meshes) cannot contain needle-like elements. If the elements are triangles, no angle can tend to zero, the so-called “minimum angle condition”. This condition is known not to be necessary for the convergence of the finite element interpolant in $H^1(\Omega)$, the necessary one being that no angle in the triangulation tend to π (the so-called “maximum angle condition”).

2.4.2 From local to global

The local estimates already obtained can be turned global by simply collecting the contributions from all elements in the mesh.

Consider the estimate of Thm. 2.7(a), to begin with. One can build an $L^\infty(\Omega)$ as follows:

$$\|u - \mathcal{I}_h u\|_{L^\infty(\Omega)} = \max_K \|u - \mathcal{I}_K u\|_{L^\infty(K)} \leq \frac{d^2}{2} \max_K \{h_K^2 \|D^2 u\|_{L^\infty(K)}\} \leq \frac{d^2}{2} h^2 \|D^2 u\|_{L^\infty(\Omega)}$$

which holds without any assumption on the mesh.

Similar estimates based on local to global reasonings are left as exercises.

Exo. 2.8 Starting from Thm. 2.7(b), prove that

$$\|\nabla u - \nabla(\mathcal{I}_h u)\|_{L^\infty(\Omega)} \leq \frac{(d+1)d^2 S}{2} h \|D^2 u\|_{L^\infty(\Omega)}$$

where S is the shape-regularity constant of the mesh. Notice that it is necessary that $\nabla(\mathcal{I}_h u)$ belongs to $L^\infty(\Omega)$, which requires a conforming mesh.

Exo. 2.9 Using (2.30) prove that

$$\|u - \mathcal{I}_h u\|_{L^2(\Omega)} \leq (d+1) \sqrt{|\Omega|} h \|\nabla u\|_{L^\infty(\Omega)} \quad (2.37)$$

Exo. 2.10 Starting from (2.32) prove that, if the family of (conforming) meshes is shape-regular and the function u smooth, then

$$\|u - \mathcal{I}_h u\|_{H^1(\Omega)} \leq C S h^k \|D^{k+1} u\|_{L^2(\Omega)} \quad (2.38)$$

where S is the shape-regularity constant of the mesh.

Exo. 2.11 Assume that there exists a straight line Γ (or planar surface in 3D) in the domain Ω , at which there is a sudden change in material properties. As a consequence, $u \in H^2(\Omega \setminus \Gamma) \cap C^0(\Omega)$, but $u \notin H^2(\Omega)$. Discuss the interpolation estimate for such a function u , showing the advantages of using an “interface-fitting mesh”; i.e., a mesh such that Γ coincides with inter-element boundaries and thus does not cut any element.

2.4.3 Global estimate

Let us state a global estimate more general than the one we have been building up to now.

Theorem 2.12 *Let \mathcal{T}_h , $h > 0$, be a family of shape-regular meshes of a domain $\Omega \subset \mathbb{R}^n$. Let $(\widehat{K}, \widehat{P}, \widehat{\Sigma})$ be the (Lagrange) reference element of the mesh, all the mappings $F_K : \widehat{K} \rightarrow K$ being affine. Let \mathcal{I}_h be the global interpolation operator corresponding to \mathcal{T}_h . Assume further that $\mathbb{P}_k \subset \widehat{P}$ (i.e.; that the finite elements are “of degree k ”). Then, for each $1 \leq p < +\infty$, and for each $0 \leq \ell \leq k$, there exists C such that for all h and all $v \in W^{\ell+1,p}(\Omega)$,*

$$\|v - \mathcal{I}_h v\|_{L^p(\Omega)} + \sum_{m=1}^{\ell+1} h^m \left(\sum_{K \in \mathcal{T}_h} |v - \mathcal{I}_h v|_{W^{m,p}(K)}^p \right)^{\frac{1}{p}} \leq C h^{\ell+1} |v|_{W^{\ell+1,p}(\Omega)} \quad (2.39)$$

If $p = +\infty$,

$$\|v - \mathcal{I}_h v\|_{L^\infty(\Omega)} + \sum_{m=1}^{\ell+1} h^m \left(\max_{K \in \mathcal{T}_h} |v - \mathcal{I}_h v|_{W^{m,\infty}(K)}^p \right)^{\frac{1}{p}} \leq C h^{\ell+1} |v|_{W^{\ell+1,\infty}(\Omega)} \quad (2.40)$$

Proof. See Ern-Guermond [?], p. 61. \square

Notice that the previous theorem holds not just for simplicial elements but also for affine-equivalent quadrilaterals, hexahedra, etc.

Exo. 2.12 *Deduce from the theorem that, for P_k and Q_k elements,*

$$\|v - \mathcal{I}_h v\|_{H^1(\Omega)} \leq C h^k, \quad \|v - \mathcal{I}_h v\|_{L^2(\Omega)} \leq C h^{k+1}$$

and explain on what quantities depend the constant C .

The previous theorem establishes, in particular, that the family of spaces $\{W_h\}$ satisfies the **approximation property**.

Prop. 2.13 *For any $v \in L^p(\Omega)$, $p < +\infty$,*

$$\lim_{h \rightarrow 0} \left(\inf_{v_h \in V_h} \|v - v_h\|_{L^p(\Omega)} \right) = 0 \quad (2.41)$$

Exo. 2.13 *Prove the previous proposition.* Hint: One cannot interpolate a generic function in $L^p(\Omega)$ because it is not continuous. Fortunately, **smooth functions are dense in $L^p(\Omega)$ for all $p < +\infty$** , so that for any $\epsilon > 0$ one can find $v^\epsilon \in H^2(\Omega)$ such that $\|v - v^\epsilon\|_{L^p(\Omega)} < \epsilon$. The interpolant $\mathcal{I}_h v^\epsilon$ is well defined and Theorem 2.12 can be applied.

2.5 Inverse inequalities

Inverse inequalities are often useful in the convergence analysis of finite element methods. They provide bounds on operators that are **unbounded** in $H^m(\Omega)$, with $m > 0$, but **bounded** in V_h due to its finite-dimensionality. Intuitively, in a shape-regular mesh for a derivative $\partial u_h / \partial x_i$ to be “very large” the nodal values of the u_h must also be “very large”.

Let $(\hat{K}, \hat{P}, \hat{\Sigma})$ be the “reference” or “master” element. Let K be an element that is affine-equivalent to \hat{K} , as defined before, with $F_K : \hat{K} \rightarrow K$ the corresponding linear mapping:

$$F_K(x) = A_K x + b_K .$$

In this section we only consider finite elements for which

$$T_K v(\hat{x}) = v(F_K(\hat{x})) ,$$

such as Lagrange finite elements. In such a setting, we have

Lemma 2.14

(a)

$$|\det A_K| = \frac{|K|}{|\hat{K}|}, \quad \|A_K\| \leq \frac{h_K}{\rho_{\hat{K}}}, \quad \|A_K^{-1}\| \leq \frac{h_{\hat{K}}}{\rho_K}$$

(b) *There exists C , depending on s and p but independent of K , such that for all $v \in W^{s,p}(K)$,*

$$|\hat{v}|_{W^{s,p}(\hat{K})} \leq C \|A_K\|^s |\det A_K|^{-\frac{1}{p}} |v|_{W^{s,p}(K)} \quad (2.42)$$

$$|v|_{W^{s,p}(K)} \leq C \|A_K^{-1}\|^s |\det A_K|^{\frac{1}{p}} |\hat{v}|_{W^{s,p}(\hat{K})} \quad (2.43)$$

Proof. See, e.g., Ciarlet [?], p. 122. \square

Let us show how to take advantage of this result to prove some simple estimates.

Prop. 2.15 *There exists $C > 0$, independent of K , such that*

$$\|\nabla v_h\|_{L^2(K)} \leq \frac{C}{\rho_K} \|v_h\|_{L^2(K)} \quad (2.44)$$

for any $v_h \in P_K$.

Proof. This proof uses the so-called *scaling* argument. From (2.43) we have, taking $s = 1$ and $p = 2$,

$$\|\nabla v_h\|_{L^2(K)} \leq C \|A_K^{-1}\| |\det A_K|^{\frac{1}{2}} \|\nabla \widehat{v}_h\|_{L^2(\widehat{K})} \quad (2.45)$$

Now let us show that there exists a constant \widehat{C} such that

$$\|\nabla \widehat{v}_h\|_{L^2(\widehat{K})} \leq \widehat{C} \|\widehat{v}_h\|_{L^2(\widehat{K})} \quad (2.46)$$

For this, consider the set $\mathcal{S} = \{w \in P_K \mid \|\widehat{w}\|_{L^2(\widehat{K})} = 1\}$, which is bounded and closed in the finite-dimensional space P_K . Let \widehat{C} be the **maximum** that the **continuous** function $\|\nabla \widehat{w}\|_{L^2(\widehat{K})}$ attains in \mathcal{S} .

Then, denoting by

$$\widehat{z}_h = \frac{1}{\|\widehat{v}_h\|_{L^2(\widehat{K})}} \widehat{v}_h$$

and noticing that $\widehat{z}_h \in \mathcal{S}$, we have that

$$\|\nabla \widehat{z}_h\|_{L^2(\widehat{K})} \leq \widehat{C}$$

and thus (2.46) is proved. Inserting it into (2.45) and using (2.42) one gets

$$\|\nabla v_h\|_{L^2(K)} \leq C \widehat{C} \|A_K^{-1}\| |\det A_K|^{\frac{1}{2}} \|\widehat{v}_h\|_{L^2(\widehat{K})} \leq C^2 \widehat{C} \|A_K^{-1}\| |\det A_K|^{\frac{1}{2}} |\det A_K|^{-\frac{1}{2}} \|v_h\|_{L^2(K)} \leq$$

$$\leq \frac{(C^2 \widehat{C} h_{\widehat{K}})}{\rho_K} \|v_h\|_{L^2(K)}$$

and the proof ends noticing that the product inside the parentheses is a constant independent of K and v_h . \square

Notice that there does **not** exist a constant C that makes

$$\|\nabla v\|_{L^2(K)} \leq \frac{C}{\rho_K} \|v\|_{L^2(K)} \quad (2.47)$$

in the **infinite dimensional case**, i.e., for any v in $H^1(K)$.

Exo. 2.14 *Let K be the unit interval $(0, 1)$ in 1D. Build a sequence $\{\varphi_n\}$ of functions such that $\|\varphi_n\|_{L^2(K)} = 1$ and $\|\nabla \varphi_n\|_{L^2(K)} = n$.*

Argue that the existence of such a sequence is a counterexample to (2.47).

With a scaling argument one can prove the following discrete trace estimate.

Prop. 2.16 *There exists $C > 0$, independent of K , such that*

$$\|v_h\|_{L^2(F)} \leq C h_K^{-\frac{1}{2}} \|v_h\|_{L^2(K)} \quad \forall v_h \in P_K \quad (2.48)$$

where F is an edge (face in 3D) of K .

The proof is left as an optional exercise. Notice that, again, there is no chance of (2.48) holding for all v in an infinite-dimensional space, such as $C^\infty(K)$ for example (build a sequence that shows this!).

Several other inverse inequalities can be extracted as particular cases of the following theorem (see, e.g., [?] p. 75).

Theorem 2.17 *Let \mathcal{T}_h be a shape-regular family of meshes in $\Omega \subset \mathbb{R}^d$. Then, for $0 \leq m \leq \ell$ and $1 \leq p, q \leq \infty$, there exists a constant C such that, for all $h > 0$ and all $K \in \mathcal{T}_h$,*

$$\|v\|_{W^{\ell,p}(K)} \leq C h_K^{m-\ell+d(\frac{1}{p}-\frac{1}{q})} \|v\|_{W^{m,q}(K)} \quad (2.49)$$

for all $v \in P_K$.

This local estimate, to be made global, puts the restriction on the family of meshes that, as $h \rightarrow 0$ the diameter ratio between the largest and smaller h_K in \mathcal{T}_h remain bounded.

Def. 2.18 *A family of meshes $\{\mathcal{T}_h\}_{h>0}$ is said to be **quasi-uniform** if it is shape-regular and there exists c such that*

$$\forall h, \quad \forall K \in \mathcal{T}_h, \quad h_K \geq ch \quad (2.50)$$

Exo. 2.15 *Does the quasi-uniformity of the mesh imply the existence of $C > 0$ such that*

$$\|\nabla v_h\|_{L^2(\Omega)} \leq C h^{-1} \|v_h\|_{L^2(\Omega)} \quad \forall v_h \in V_h ? \quad (2.51)$$

Exo. 2.16 *Does the quasi-uniformity of the mesh imply the existence of $C > 0$ such that*

$$\|v_h\|_{L^2(\partial\Omega)} \leq C h^{-\frac{1}{2}} \|v_h\|_{L^2(\Omega)} \quad \forall v_h \in V_h ? \quad (2.52)$$

Let us now give a try at the easy part of the proof of Theorem 2.31.

The point of departure is the following proposition, in which the carets are inserted to remind that it is to be applied on the master element:

Prop. 2.19 *Let $\mathbb{P}_k \subset \widehat{P}$, $k \geq 1$, then there exists $\widehat{C} > 0$ such that*

$$\|\hat{u} - \mathcal{I}_{\widehat{K}}\hat{u}\|_{L^2(\widehat{K})} + \|\nabla\hat{u} - \nabla\mathcal{I}_{\widehat{K}}\hat{u}\|_{L^2(\widehat{K})} \leq \widehat{C} \|D^{k+1}\hat{u}\|_{L^2(\widehat{K})} \quad \forall u \in H^{k+1}(\widehat{K}) . \quad (2.53)$$

Exo. 2.17 *Taking Proposition 2.19 as established, prove Theorem 2.31. The strategy is a scaling argument analogous to that used in the proof of Prop. 2.15.*

Now, how to prove Prop. 2.19? Leaving the details to be read from the literature, let us just put forward the main conceptual ingredient:

Theorem 2.20 (Bramble-Hilbert lemma) *Let $F : H^{k+1}(\omega) \rightarrow \mathbb{R}$ be a continuous linear functional, satisfying*

$$F(p) = 0 , \quad \forall p \in \mathbb{P}_k . \quad (2.54)$$

Assuming $\omega \subset \mathbb{R}^d$ to be convex and bounded, with Lipschitz boundary, there exists $C(\omega) > 0$ such that

$$|F(v)| \leq C(\omega) \|D^{k+1}v\|_{L^2(\omega)} , \quad \forall v \in H^{k+1}(\omega) . \quad (2.55)$$

2.6 Interpolation in $H(\mathbf{div}, \Omega)$

The Raviart-Thomas element introduced in Exo. 2.4 has **constant normal component** on each face of K , leading to

$$\sigma_i(v_h|_K) = \text{meas}(F_i) (v_h \cdot \tilde{n})(F_i) . \quad (2.56)$$

Because $(v_h \cdot \tilde{n})(F_i)$ is single-valued for the two elements sharing face F_i , the global space W_h generated by RT_0 elements is a subspace of $H(\mathbf{div}, \Omega)$.

The interpolant $\mathcal{I}_h^{RT} : H(\mathbf{div}, \Omega) \rightarrow W_h$ is built using the local-to-global construction as before.

With the same ingredients as before (Bramble-Hilbert lemma, scaling arguments) it is possible to prove the following approximation result:

Theorem 2.21 *Let \mathcal{T}_h be a shape-regular family of triangulations. There exists $c > 0$ such that, for all h and for all $v \in H^1(\Omega)^d$ with $\nabla \cdot v \in H^1(\Omega)$,*

$$\|v - \mathcal{I}_h^{RT} v\|_{L^2(\Omega)} + \|\nabla \cdot (v - \mathcal{I}_h^{RT} v)\|_{L^2(\Omega)} \leq ch \left(\|v - \mathcal{I}_h^{RT} v\|_{H^1(\Omega)} + \|\nabla \cdot (v - \mathcal{I}_h^{RT} v)\|_{H^1(\Omega)} \right) \quad (2.57)$$

2.7 Interpolation of non-smooth functions

As already mentioned, Lagrange interpolation is not defined for arbitrary functions in $L^p(\Omega)$, not even for $H^1(\Omega)$ if $d > 1$. In applications that will be discussed later on, it is important that there exists an interpolation operator $\bar{\mathcal{I}}_h : H^1(\Omega) \rightarrow W_h$ with some useful properties:

- **Stability:** There exists $c > 0$ such that

$$\forall h, \forall v \in L^2(\Omega), \quad \|\bar{\mathcal{I}}_h v\|_{L^2(\Omega)} \leq c \|v\|_{L^2(\Omega)}. \quad (2.58)$$

$$\forall h, \forall v \in H^1(\Omega), \quad \|\bar{\mathcal{I}}_h v\|_{H^1(\Omega)} \leq c \|v\|_{H^1(\Omega)}. \quad (2.59)$$

- **Approximation:** There exists $c > 0$ such that

$$\forall h, \forall K \in \mathcal{T}_h, \forall v \in H^1(\omega_K), \quad \|v - \bar{\mathcal{I}}_h v\|_{L^2(K)} \leq ch \|v\|_{H^1(\omega_K)} \quad (2.60)$$

where ω_K consists of all elements sharing at least a node with K .

- **Preservation of boundary conditions:** If $v|_{\partial\Omega} = 0$, then $(\bar{\mathcal{I}}_h v)|_{\partial\Omega} = 0$.
- **Being a projection:** $\bar{\mathcal{I}}_h v = v$ for all $v \in W_h$.

The **Clément interpolation** operator satisfies the first two properties, while the **Scott-Zhang interpolation** operator satisfies all four.

Exo. 2.18 Read the construction of the Clément and Scott-Zhang interpolation operators, for example in Ern & Guermond, p. 68-71.

3 Galerkin treatment of elliptic second-order problems

3.1 The continuous problem

We consider the following problem:

$$-\operatorname{div}(K\nabla u) + \beta \cdot \nabla u + \sigma u = f \quad \text{in } \Omega \quad (3.1)$$

$$u = g \quad \text{on } \Gamma_D \quad (3.2)$$

$$(K\nabla u) \cdot \mathbf{n} = H \quad \text{on } \Gamma_N \quad (3.3)$$

where Γ_D and Γ_N are disjoint parts of $\partial\Omega$, and $\overline{\Gamma_D} \cup \overline{\Gamma_N} = \partial\Omega$.

Notice that, since $K(x)$ is a $n \times n$ symmetric matrix and $\beta(x)$ is an n -vector, the problem above is a general second-order partial differential equation.

Integrating formally by parts we get

$$\int_{\Omega} (\nabla v \cdot (K\nabla u) + v \beta \cdot \nabla u + \sigma uv) \, d\Omega = \int_{\Omega} f v \, d\Omega + \int_{\partial\Omega} v \mathbf{n} \cdot (K\nabla u) \, d\Gamma$$

We thus consider the bilinear form

$$a(u, v) = \int_{\Omega} (\nabla v \cdot (K\nabla u) + v \beta \cdot \nabla u + \sigma uv) \, d\Omega \quad (3.4)$$

Prop. 3.1 *If $K \in (L^\infty(\Omega))^{n \times n}$, $\beta \in (L^\infty(\Omega))^n$ and $\sigma \in L^\infty(\Omega)$, then $a(\cdot, \cdot)$ is continuous on $H^1(\Omega)$.*

Exo. 3.1 *Prove the proposition.*

It is clear that, for the problem to admit a solution, the data g and Γ_D must be regular enough for a function $u_g \in H^1(\Omega)$ to exist satisfying $u_g = g$ on Γ_D . Such a function is called a “lifting” function, and if it exists one says that g belongs to a “trace space”.

We now change the unknown to $w = u - u_g$, so that

$$a(w, v) = \int_{\Omega} f v \, d\Omega + \int_{\partial\Omega} v \mathbf{n} \cdot (K \nabla u) \, d\Gamma - a(u_g, v)$$

and $w = 0$ on Γ_D . This leads us to consider the following problem: *Find $w \in H_{D_0}^1(\Omega)$ such that*

$$a(w, v) = \int_{\Omega} f v \, d\Omega + \int_{\Gamma_N} H v \, d\Gamma - a(u_g, v) \stackrel{\text{def}}{=} \ell(v) \quad (3.5)$$

where $H_{D_0}^1 = \{v \in H^1(\Omega), v = 0 \text{ on } \Gamma_D\}$.

Prop. 3.2 *Assume the data f, g, H, Γ_N and Γ_D are regular enough for the right-hand side of (3.5) to be a continuous linear functional on $H_{D_0}^1(\Omega)$. Assume further that the hypotheses of Prop. 3.1 hold, and that*

$$\operatorname{div} \beta \in L^\infty(\Omega), \quad \beta(x) \cdot n(x) > 0 \quad \text{a.e. on } \Gamma_N \quad (3.6)$$

$$\xi \cdot (K(x)\xi) \geq K_0 |\xi|^2 \quad \forall \xi \in \mathbb{R}^n; \text{ a.e. in } \Omega \quad (3.7)$$

$$\sigma(x) - \frac{1}{2} \operatorname{div} \beta(x) \geq s_{\min} \quad \text{a.e. in } \Omega \quad (3.8)$$

where K_0 and s_{\min} are strictly positive constants. Then (3.5) is well-posed.

Proof. Notice first that $H_{D0}^1(\Omega)$ is a closed subspace of $H^1(\Omega)$. To see this, consider the applications $\gamma_0 : H^1(\Omega) \rightarrow L^2(\partial\Omega)$ (the boundary trace operator, which is continuous as proved for example in Adams, Brenner-Scott, etc.) and $r_D : L^2(\partial\Omega) \rightarrow L^2(\Gamma_D)$, the restriction to Γ_D of a function in $L^2(\Omega)$, which is also continuous. The value of any function $f \in H^1(\Omega)$ on Γ_D is, then, $\gamma_{0D}(f) = r_D(\gamma_0(f))$. The subspace $H_{D0}^1(\Omega)$ is the pre-image of zero by γ_{0D} , and is thus closed.

To conclude the proof, it remains to show that $a(\cdot, \cdot)$ is weakly coercive. In fact, a direct calculation shows that $a(\cdot, \cdot)$ is strongly coercive and thus Lax-Milgram lemma guarantees well-posedness. \square

The condition

$$\xi \cdot (K(x)\xi) \geq K_0 |\xi|^2 > 0, \quad \forall \xi \in \mathbb{R}^n; \text{ a.e. in } \Omega$$

is essential to the previous well-posedness result, as it applies only for *elliptic* second-order PDEs (not hyperbolic, not parabolic). The condition $s_{\min} > 0$ is not essential, in the sense that if $s_{\min} \leq 0$ what may happen is that the *homogeneous* problem defined by $f = g = H = 0$ admits non-trivial solutions. It may also happen that for certain data the solution does not exist, in much the same way as a linear system

$$\underline{\underline{A}} \underline{x} = \underline{b}$$

with $\det(\underline{\underline{A}}) = 0$ either does not have a solution, or has infinitely many (the solution is determined only up to the addition of an arbitrary element of $\text{Ker}(\underline{\underline{A}})$).

Exa. 3.3 *The simplest and very important case that is not covered by Prop. 3.2 is the purely diffusive problem with Neumann data, corresponding to*

$$\beta = 0 \text{ (no convection)}, \quad \sigma = 0 \text{ (no reaction)}, \quad \Gamma_N = \partial\Omega \text{ (no Dirichlet boundary)}. \quad (3.9)$$

The differential formulation is

$$- \text{div}(K\nabla u) = f \quad \text{in } \Omega, \quad (K\nabla u) \cdot \mathbf{n} = H \quad \text{on } \partial\Omega \quad (3.10)$$

which only admits a solution if

$$\int_{\Omega} f + \int_{\partial\Omega} H = 0$$

and, in this case, the solution is determined up to an additive constant. Notice that the constant functions are indeed solutions of the homogeneous problem ($f = H = 0$), and in fact the only solutions if Ω is connected.

Exo. 3.2 Show that under the hypotheses of Prop. 3.2 the bilinear form $a(\cdot, \cdot)$ is indeed strongly coercive (as claimed) and provide an estimate of the coercivity constant α .

Let now $H_{Dg}^1(\Omega) = \{v \in H^1(\Omega); v = g \text{ a.e. on } \Gamma_D\}$. Setting $u = u_g + w$ it is clear that u solves the following problem: Find $u \in H_{Dg}^1(\Omega)$ such that

$$a(u, v) = \int_{\Omega} f v \, d\Omega + \int_{\Gamma_N} H v \, d\Gamma \quad (3.11)$$

for all $v \in H_{D0}^1(\Omega)$.

Further, if u belongs to $H^2(\Omega)$ integration by parts shows that the partial differential equation holds almost everywhere in Ω and that the Neumann boundary condition is satisfied on Γ_N .

Notice that the Neumann boundary condition enters the right-hand side of (3.11), it is a *natural* condition for this formulation, while the Dirichlet condition has to be imposed to the space in which the solution is sought, it is an *essential* boundary condition. One could wonder whether the Neumann boundary condition could also be imposed as an essential condition: The answer is that the set of functions in $H^1(\Omega)$ which satisfy $\mathbf{n} \cdot (K\nabla u) = H$ on Γ_N is *not* closed in $H^1(\Omega)$, implying that the tools we use to prove existence (the Banach and Hahn-Banach theorems in the general case, the Lax-Milgram lemma in the strongly coercive, Hilbertian case) do not apply.

Exo. 3.3 Let $\Omega = (0, 1)$. Let $\varphi(x) = x$. Show a sequence $\{\varphi_n\} \subset H^1(\Omega)$ such that $\varphi'_n(0) = 0$ for all n and such that $\varphi_n \rightarrow \varphi$ strongly in $H^1(\Omega)$.

Hint: For $1/n = \epsilon > 0$ consider the “trimmed” function

$$T_\epsilon \varphi(x) = \begin{cases} \varphi(\epsilon) & \text{if } x < \epsilon \\ \varphi(x) & \text{if } x \geq \epsilon \end{cases}$$

3.2 Ritz-Galerkin approximation

Let $V_h(\Omega)$ be a finite element space contained in $H^1(\Omega)$, and let $V_{h0}(\Omega)$ be the subspace of $V_h(\Omega)$ obtained by putting to zero all degrees of freedom corresponding to values on Γ_D . Analogously, $V_{hg}(\Omega)$ is defined as the (linear) subset of $V_h(\Omega)$ consisting of functions that coincide with some given interpolation $I_h g$ of g on Γ_D . The Ritz-Galerkin approximation of u in $V_h(\Omega)$ then solves:

Find $u_h \in V_{hg}(\Omega)$ such that

$$a(u_h, v_h) = \int_{\Omega} f v_h \, d\Omega + \int_{\partial\Omega} H v_h \, d\Gamma \quad (3.12)$$

for all $v_h \in V_{h0}(\Omega)$.

Applying Lax-Milgram lemma to the discrete problem immediately implies that it is well-posed. By Céa’s lemma (Lemma 1.39),

$$\|u - u_h\|_1 \leq \frac{N_a}{\alpha} \inf_{v_h \in V_{hg}(\Omega)} \|u - v_h\|_1 \leq \frac{N_a}{\alpha} \|u - \mathcal{I}_h u\|_1$$

Thus, if the local space P_K on each element K of the mesh \mathcal{T}_h contains all polynomials up to degree k and the solution is smooth enough,

$$\|u - u_h\|_1 \leq Ch^k |u|_{k+1}$$

3.3 Aubin-Nitsche's duality argument

The error bound in the $H^1(\Omega)$ -norm, as shown before, is naturally obtained in the Ritz-Galerkin formulation of second-order PDEs. A first estimate in the $L^2(\Omega)$ -norm follows from the continuous injection of $H^1(\Omega)$ into $L^2(\Omega)$, yielding

$$\|u - u_h\|_0 \leq Ch^k |u|_{k+1}$$

This estimate, however, is not optimal, since the interpolant of u (with u smooth) approximates u with order h^{k+1} in the $L^2(\Omega)$ -norm. It is possible to obtain optimal-order estimates using a duality argument. Let us show how it works in the simpler case $\beta = 0$, $g = 0$, $\Gamma_D = \partial\Omega$. Let

$$\mathcal{L}u = -\operatorname{div}(K\nabla u) + \sigma u$$

and assume that the domain is regular enough for \mathcal{L} to have a *smoothing property*, namely that the continuous problem

$$\mathcal{L}w = \mathcal{F}, \quad w = 0 \quad \text{on } \partial\Omega$$

satisfies

$$\|w\|_{H^2(\Omega)} \leq C_s \|\mathcal{F}\|_{L^2(\Omega)} \tag{3.13}$$

This latter inequality is sometimes called a *regularity estimate*.

Exo. 3.4 *Prove the smoothing property in 1D. More specifically, consider the problem*

$$-(ku')' + \sigma u = f \quad \text{in } \Omega = (0, 1) \tag{3.14}$$

with $u(0) = u(1) = 0$, $k, \sigma \in L^\infty(\Omega)$ satisfying $k(x) \geq \gamma > 0$ for all x and $\sigma(x) \geq 0$ for all x . Further, assume that $k' \in L^\infty(\Omega)$, $f \in L^2(\Omega)$. Notice that $k'(x)$ must be bounded. Show that then there exists $C > 0$ such that $\|u''\|_{L^2(\Omega)} \leq C \|f\|_{L^2(\Omega)}$ and provide an estimate for C . Show how this implies (3.13).

Remark 3.4 *The smoothing property (3.13) holds in 2D/3D if the boundary is very regular, of class C^2 , or if it is a convex polygon/polyhedron.*

Prop. 3.5 *Under the above hypotheses, there exists $C > 0$ such that*

$$\|u - u_h\|_0 \leq Ch\|u - u_h\|_1 \quad (3.15)$$

Proof. Let w be the unique solution of

$$\mathcal{L}w = u - u_h, \quad w = 0 \quad \text{on } \partial\Omega$$

where we have used the error $e = u - u_h$ as source term. The corresponding variational formulation is

$$a(w, v) = (e, v)_0 \quad \forall v \in H_0^1(\Omega)$$

Taking $v = e$ we see that $a(w, e) = \|e\|_0^2$, but also, since the bilinear form is symmetric (otherwise one needs a smoothing property for the adjoint differential operator, but the proof is essentially the same),

$$a(w, e) = a(e, w) = a(u - u_h, w) = a(u - u_h, w - \mathcal{I}_h w)$$

where we have introduced the interpolant of w and used the “orthogonality” property of the Galerkin approximation ($a(u - u_h, v_h) = 0$ for all v_h). Finally

$$\|u - u_h\|_0^2 = a(e, w - \mathcal{I}_h w) \leq N_a \|e\|_1 \|w - \mathcal{I}_h w\|_1 \leq N_a \|e\|_1 h \|w\|_2$$

where the last inequality follows from an interpolation estimate for w . Combining with (3.13),

$$\|u - u_h\|_0^2 \leq C_s N_a h \|e\|_1 \|e\|_0$$

□

Exo. 3.5 *Let $F(v) = \int_{\Omega} \psi(x) v(x) d\Omega$, where ψ is a function in $L^2(\Omega)$. For example, if $\psi = 1$ then $F(v)$ is simply the integral of v . How does $F(u_h)$ converge to $F(u)$ when V_h contains all piecewise polynomials*

of degree k and $\|u - u_h\|_1 \leq C h^k$?

Hint: Use a variant of Nitsche's trick. Let w be the solution of

$$a(w, v) = F(v) \quad \forall v \in V = H_0^1(\Omega)$$

which is the weak form of

$$\mathcal{L}w = \psi \quad \text{in } \Omega, \quad w = 0 \quad \text{on } \partial\Omega$$

so that, from the smoothing property, $\|w\|_{H^2(\Omega)} \leq C \|\psi\|_{L^2(\Omega)}$. Then use the following calculation

$$F(u - u_h) = a(w, u - u_h) = a(w - \mathcal{I}_h w, u - u_h) \leq N_a \|w - \mathcal{I}_h w\|_1 \|u - u_h\|_1$$

to prove that, if ψ is smooth (at least as smooth as f), then $|F(u) - F(u_h)| \leq \tilde{C} h^{2k}$.

Another question: What is the expected order of convergence for $F(u) = \int_\omega u \, d\Omega$, with ω a region of the domain? (Answer: h^{k+1} , why?).

3.4 The case $s_{\min} = 0$. Poincaré inequality.

In the case $s_{\min} = 0$ we have to prove strong coercivity without counting on the reaction term, so that we start from the estimate

$$a(v, v) \geq \int_\Omega \nabla v \cdot (K \nabla v) \, d\Omega \quad \forall v \in H_{D_0}^1(\Omega)$$

which in turn implies

$$a(v, v) \geq K_0 \int_\Omega |\nabla v|^2 \, d\Omega = K_0 |v|_1^2$$

Essentially, we need an estimate of the form $|v|_1 \geq c \|v\|_1$ for some $c > 0$. This is provided by Poincaré-Friedrichs inequality:

Lemma 3.6 (Poincaré-Friedrichs inequality) *In a connected bounded domain, if $\text{meas}(\Gamma_D) > 0$ then there exists a constant $c_P > 0$ such that $\|\nabla v\|_0 \geq c_P \|v\|_0$ for all $v \in H_{D_0}^1(\Omega)$.*

Proof. We will prove it in the case $\Gamma_D = \partial\Omega$. We show it first for $\varphi \in \mathcal{D}(\Omega)$ and then extend it to $H_0^1(\Omega)$ by a density argument. We consider φ extended by zero to \mathbb{R}^n and assume that the domain is contained in the strip $a \leq x_1 \leq b$ (in other words, $a \leq x_1 \leq b$ for all $x \in \Omega$). Then, since

$$\varphi(x_1, x_2, \dots, x_n) = \int_a^{x_1} \frac{\partial \varphi}{\partial x_1}(t, x_2, \dots, x_n) dt$$

we have, using Cauchy-Schwarz inequality,

$$\varphi^2(x_1, x_2, \dots, x_n) \leq |x_1 - a| \int_a^{x_1} \left| \frac{\partial \varphi}{\partial x_1}(t, x_2, \dots, x_n) \right|^2 dt$$

integration over x_2 to x_n gives

$$\int \varphi^2 dx_2 \dots dx_n \leq |x_1 - a| \int_a^{x_1} \dots \int \left| \frac{\partial \varphi}{\partial x_1} \right|^2 dt dx_2 \dots dx_n \leq |x_1 - a| \int_{\Omega} \left| \frac{\partial \varphi}{\partial x_1} \right|^2 d\Omega$$

A final integration over x_1 yields

$$\int_{\Omega} \varphi^2 d\Omega \leq \frac{(b-a)^2}{2} \int_{\Omega} \left| \frac{\partial \varphi}{\partial x_1} \right|^2 d\Omega$$

proving that $c_P \geq \frac{\sqrt{2}}{b-a}$. Now we consider $v \in H_0^1(\Omega)$ and $\varphi_n \rightarrow v$, then

$$\|v\|_0 \leq \|\varphi_n\|_0 + \|v - \varphi_n\|_0 \leq \frac{1}{c_P} \|\nabla \varphi_n\|_0 + \|v - \varphi_n\|_0 \leq$$

$$\frac{1}{c_P} \|\nabla v\|_0 + \|v - \varphi_n\|_0 + \frac{1}{c_P} \|\nabla v - \nabla \varphi_n\|_0 \leq \frac{1}{c_P} \|\nabla v\|_0 + \min \left\{ 1, \frac{1}{c_P} \right\} \|v - \varphi_n\|_1$$

and since $\|v - \varphi_n\|_1$ can be made arbitrarily small, the claim is proved. \square

Remark 3.7 *Using Poincaré-Friedrichs inequality, it is easily shown that the bilinear form*

$$a(v, w) = \int_{\Omega} [\nabla v \cdot (K \nabla w) + v \beta \cdot \nabla w + \sigma v w] \, d\Omega \quad (3.16)$$

is strongly coercive in $H_{D_0}^1(\Omega)$ whenever $\text{meas}(\Gamma_D) > 0$, $\beta(x) \cdot \mathbf{n}(x) > 0$ a.e. on Γ_N , $K_0 > 0$ and $s_{\min} \geq 0$.

Exo. 3.6 *Prove the previous remark in detail.*

4 Finite elements for linear elasticity

4.1 Introduction and differential formulation

We recall the usual notations for the Cauchy stress tensor $\boldsymbol{\sigma}$ and the linearized strain tensor

$$\boldsymbol{\epsilon}(u) = \frac{1}{2} (\nabla u + \nabla u^T) \quad (4.1)$$

where u in this case is a *vector field* corresponding to the *displacement* of the body. We also recall the elastic constitutive law for small deformations,

$$\boldsymbol{\sigma} = \lambda \operatorname{tr}(\boldsymbol{\epsilon}(u)) \mathbf{I} + 2\mu \boldsymbol{\epsilon}(u) = \lambda \operatorname{div} u \mathbf{I} + \mu (\nabla u + \nabla u^T) \quad (4.2)$$

where λ and μ are the Lamé coefficients, which in general depend on the point x and by thermodynamic reasons are constrained to satisfy, for almost all x ,

$$\mu(x) > 0; \quad \lambda(x) + \frac{2}{3} \mu(x) \geq 0 \quad (4.3)$$

Differential Formulation: The governing equation follows from the static equilibrium balance, which reads

$$\operatorname{div} \boldsymbol{\sigma} + f = 0 \quad (4.4)$$

where f is a vector field of applied forces. Replacing the expression of $\boldsymbol{\sigma}$ in terms of u one obtains an equation for the displacement field. This problem admits both Dirichlet and Neumann boundary conditions on u :

$$u = g \quad \text{on } \Gamma_D; \quad \boldsymbol{\sigma} \cdot \mathbf{n} = \mathcal{F} \quad \text{on } \Gamma_N \quad (4.5)$$

where \mathcal{F} is a field of surface forces applied on Γ_N , $\Gamma_N \cap \Gamma_D = \emptyset$ and $\overline{\Gamma_N \cup \Gamma_D} = \partial\Omega$. The domain Ω corresponds to the region of space occupied by the body under consideration, both before and after the application of the forces since just problems with *small displacements* are being considered.

Exo. 4.1 Let u_1, u_2 be the components of u in a planar elasticity case in which the domain is the unit square. The boundary conditions are: zero displacement on the bottom boundary ($x_2 = 0$), and a normal force equal to P on the rest of $\partial\Omega$. Write down the system of two equations and two unknowns for u_1 and u_2 considering λ and μ independent of x_1 and x_2 .

Hint: Equation (4.4), written in Cartesian indices, becomes

$$\sum_{j=1}^d \partial_j \sigma_{ij} + f_i = 0 \quad \forall i = 1, \dots, d$$

and (4.2) becomes,

$$\sigma_{ij} = \lambda(\partial_1 u_1 + \partial_2 u_2) \delta_{ij} + \mu (\partial_j u_i + \partial_i u_j).$$

It remains to replace the latter into the former. For the boundary force we have that, if $\mathbf{x} = (x_1, x_2) \in \partial\Omega$ then at \mathbf{x} we have

$$(\boldsymbol{\sigma} \cdot \mathbf{n})_1 = [(\lambda + 2\mu)\partial_1 u_1 + \lambda\partial_2 u_2] n_1 + \mu (\partial_2 u_1 + \partial_1 u_2) n_2 = -P n_1$$

$$(\boldsymbol{\sigma} \cdot \mathbf{n})_2 = [\lambda\partial_1 u_1 + (\lambda + 2\mu)\partial_2 u_2] n_2 + \mu (\partial_2 u_1 + \partial_1 u_2) n_1 = -P n_2$$

As a consequence, along $x_1 = 0$ (left boundary), the boundary conditions are

$$(\lambda + 2\mu)\partial_1 u_1 + \lambda\partial_2 u_2 = -P, \quad \partial_2 u_1 + \partial_1 u_2 = 0$$

the conditions at the other boundaries are analogous.

4.2 Variational Formulation

The variational formulation of this problem can be obtained from the corresponding PDE by integration by parts. In Mechanics, however, it is considered a *fundamental principle*: The Principle of Virtual Work (or of Virtual Power)

Principle of Virtual Power: The internal virtual power of the stresses $(\int_{\Omega} \boldsymbol{\sigma} : \boldsymbol{\epsilon}(v))$ plus the virtual power of the acceleration $(\int_{\Omega} \rho a \cdot v)$ equals the virtual power of the applied forces. This holds for all virtual velocity fields, that is, all vector fields v that are kinematically admissible variations of the body motion.

$$\int_{\Omega} \boldsymbol{\sigma} : \boldsymbol{\epsilon}(v) + \int_{\Omega} \rho a \cdot v = \int_{\Omega} f \cdot v + \int_{\Gamma_N} \mathcal{F} \cdot v \quad \forall v \in \text{VAR} \quad (4.6)$$

The kinematically admissible motions must belong to

$$\text{KIN} = V_{Dg} = \{v \in [H^1(\Omega)]^n; v = g \text{ on } \Gamma_D\} \quad (4.7)$$

so that their variations must belong to

$$\text{VAR} = V_{D0} = \{v \in [H^1(\Omega)]^n; v = 0 \text{ on } \Gamma_D\} \quad (4.8)$$

The variational formulation of **linear elastostatics** then reads:

“Find $u \in V_{Dg}$ such that

$$a(u, v) = \int_{\Omega} f \cdot v \, d\Omega + \int_{\Gamma_N} \mathcal{F} \cdot v \, d\Gamma =: \ell(v) \quad (4.9)$$

for all $v \in V_{D0}$ ”, where

$$a(u, v) = \int_{\Omega} \boldsymbol{\sigma}(\mathbf{u}) : \boldsymbol{\epsilon}(v) \, d\Omega = \int_{\Omega} [\lambda \operatorname{div} u \operatorname{div} v + 2\mu \boldsymbol{\epsilon}(u) : \boldsymbol{\epsilon}(v)] \, d\Omega \quad (4.10)$$

4.3 Well-posedness and Galerkin approximation

Theorem 4.1 (Korn’s inequality) *Let Ω be a domain in \mathbb{R}^n . There exists $C_K > 0$ such that*

$$\|v\|_1 \leq C_K \|\boldsymbol{\epsilon}(v)\|_0 \quad \forall v \in H_0^1(\Omega)^n \quad (4.11)$$

It is not necessary that v be zero on the whole of $\partial\Omega$, the same result holds if $\operatorname{meas}(\Gamma_D) > 0$ (in connected domains), so that we have strong coercivity of the bilinear form on V . This gives the result below.

Theorem 4.2 *Let Ω be a regular domain on which the elasticity problem (4.9) is posed with $\operatorname{meas}(\Gamma_D) > 0$, $f \in L^2(\Omega)^n$ and $\mathcal{F} \in L^2(\Gamma_N)^n$. We assume that the Lamé coefficients are bounded and satisfy (4.3). Then there exists a unique solution u , and there exists $c > 0$ such that*

$$\|u\|_1 \leq c (\|f\|_0 + \|\mathcal{F}\|_{0, \Gamma_N}) \quad (4.12)$$

Proof. $V = V_{D0}$ is a Hilbert space, the bilinear form is continuous with

$$a(u, v) \leq c \max \{\lambda_{\max}, \mu_{\max}\} \|\nabla u\|_0 \|\nabla v\|_0$$

From Korn's inequality we also have

$$a(v, v) = \int_{\Omega} [\lambda(\operatorname{div} v)^2 + 2\mu\epsilon(v) : \epsilon(v)] \, d\Omega \geq c\mu_{\min}\|v\|_1^2$$

It only remains to apply Lax-Milgram lemma.

□

Let

$$V_h = \{v_h \in C^0(\bar{\Omega})^n, v_h|_K \in (P_K)^n, v_h = 0 \text{ on } \Gamma_D\}. \quad (4.13)$$

Since $V_h \subset V$, we have well-posedness and convergence of the discrete problem.

Prop. 4.3 *The solution $u_h \in V_{hg}$ satisfying*

$$a(u_h, v_h) = \ell(v_h) \quad \forall v_h \in V_{hg} \quad (4.14)$$

exists and is unique. It satisfies $\lim_{h \rightarrow 0} \|u - u_h\|_1 = 0$. If $u \in H^{\ell+1}(\Omega)^n$ for some $\ell \leq k$, with k such that $P_k(K) \subset P_K$, then there exists $c > 0$ such that

$$\|u - u_h\|_1 \leq ch^\ell |u|_{\ell+1} \quad (4.15)$$

Exo. 4.2 *Build an extremal formulation of the linear elasticity problem.*

Hint: Consider

$$J(w) = \int_{\Omega} \left[\frac{\lambda}{2} (\operatorname{div} w)^2 + \mu \epsilon(w) : \epsilon(w) \right] \, d\Omega - \int_{\Omega} f \cdot w \, d\Omega - \int_{\Gamma_N} \mathcal{F} \cdot w \, d\Gamma \quad (4.16)$$

where the first integral is the “strain energy” of the body. The solution u is the displacement field that minimizes J over V_{Dg} ,

$$J(u) = \inf_{w \in V_{Dg}} J(w) \quad (4.17)$$

4.4 Implementation aspects

A significant difference between the elastostatics problem and the convection-diffusion-reaction problem discussed earlier is that the elasticity unknown is a vector field.

Let $\{\mathcal{N}^j\}$ ($j = 1, \dots, M$) be the scalar basis functions associated to a mesh \mathcal{T}_h . The space V_h is now of dimension $n \times M$, as to each node j correspond n basis functions:

$$\mathbf{N}^{j,1}(x) = \mathcal{N}^j(x) \check{\mathbf{e}}^1 = (\mathcal{N}^j(x), 0) \quad \dots \quad \mathbf{N}^{j,n}(x) = \mathcal{N}^j(x) \check{\mathbf{e}}^n = (0, \mathcal{N}^j(x)) \quad (4.18)$$

where we have chosen the local basis $\{\check{\mathbf{e}}^\alpha\}$ equal to the canonical basis ($\check{\epsilon}_\beta^\alpha = \delta_{\alpha\beta}$), but any other can be chosen and sometimes is.

Exo. 4.3 *Compute the following in terms of the scalar basis $\{\mathcal{N}^j\}$:*

- $div(\mathbf{N}^{j,\alpha})$ (*Answer:* $= \partial_\alpha \mathcal{N}^j$)
- $\epsilon(\mathbf{N}^{j,\alpha})$
- $\int_K div(\mathbf{N}^{j,\alpha}) div(\mathbf{N}^{k,\beta})$
- $\int_K \epsilon(\mathbf{N}^{j,\alpha}) : \epsilon(\mathbf{N}^{k,\beta})$

5 Finite elements for mixed problems

5.1 Constraints and Lagrange multipliers

It is typical of applications of the FEM to involve *constraints* on the admissible set of solutions. Let us briefly describe some examples.

5.1.1 Incompressible elasticity

There exist elastic materials which behave as incompressible, in the sense that they preserve their volume in every deformation. Under the hypothesis of small deformations, the preservation of volume is equivalent to the deformation field having zero divergence,

$$\operatorname{div} u = 0 \quad \text{a.e. in } \Omega \quad (5.1)$$

Considering the elastic energy functional seen in the previous section (where λ is assumed independent of x for simplicity and $\|\epsilon(v)\|^2 = \epsilon(v) : \epsilon(v)$)

$$J(v) = \frac{\lambda}{2} \int_{\Omega} (\operatorname{div} v)^2 d\Omega + \int_{\Omega} \mu \|\epsilon(v)\|^2 d\Omega - \int_{\Omega} f \cdot v d\Omega - \int_{\Gamma_N} \mathcal{F} \cdot v d\Gamma \quad (5.2)$$

one can view the first term as a penalization (with coefficient λ) of the incompressibility constraint. As a consequence, totally incompressible behavior corresponds to $\lambda \rightarrow +\infty$ in theory, and to λ very large, much larger than the shear modulus μ , in practice.

For the *Primal Formulation*, which is the one we have been studying up to now, the divergence-free constraint is treated as an *essential constraint*, just like the Dirichlet constraints, and is incorporated into the set of admissible displacement fields,

$$Z_{Dg} \stackrel{\text{def}}{=} \{v \in V_{Dg} \mid \operatorname{div} v = 0 \text{ a.e. in } \Omega\} \quad (5.3)$$

Inside Z_{Dg} the first term of J becomes irrelevant, so that defining

$$\tilde{J}(v) = \int_{\Omega} \mu \|\epsilon(v)\|^2 d\Omega - \int_{\Omega} f \cdot v d\Omega - \int_{\Gamma_N} \mathcal{F} \cdot v d\Gamma, \quad (5.4)$$

we have the *Primal Extremal Formulation of incompressible elasticity*.

Primal Extremal Formulation of incompressible elasticity: Find $u \in Z_{Dg}$ that minimizes \tilde{J} over Z_{Dg} ,
i.e.,

$$\tilde{J}(u) \leq \tilde{J}(v) \quad \forall v \in Z_{Dg} \quad (5.5)$$

Defining now

$$\tilde{a}(u, v) = \int_{\Omega} 2\mu \epsilon(u) : \epsilon(v) d\Omega, \quad \text{and} \quad \ell(v) = \int_{\Omega} f \cdot v d\Omega + \int_{\Gamma_N} \mathcal{F} \cdot v d\Gamma \quad (5.6)$$

we have

$$\tilde{J}(v) = \frac{1}{2} \tilde{a}(v, v) - \ell(v) \quad (5.7)$$

and also the

Primal Variational Formulation of incompressible elasticity: Find $u \in Z_{Dg}$ such that

$$\tilde{a}(u, v) = \ell(v) \quad \forall v \in Z_{D0} \quad (5.8)$$

It can be shown that problem (5.8) is indeed well posed, so that a unique solution u exists. However, the imposition of the zero-divergence constraint on the space creates several difficulties for the finite element discretization.

It is thus convenient to replace the Primal Extremal Formulation by the following equivalent one:

Mixed Extremal Formulation of incompressible elasticity: Defining $b(\cdot, \cdot) : H^1(\Omega)^d \times L^2(\Omega) \rightarrow \mathbb{R}$ by

$$b(v, q) = \int_{\Omega} q \operatorname{div} v \, d\Omega \quad (5.9)$$

and the Lagrangian $\mathcal{L} : H^1(\Omega)^d \times L^2(\Omega) \rightarrow \mathbb{R}$ by

$$\mathcal{L}(v, q) = \tilde{J}(v) - b(v, q) = \frac{1}{2} \tilde{a}(v, v) - \ell(v) - b(v, q) , \quad (5.10)$$

problem (5.5) becomes equivalent to “Find $(u, p) \in V_{Dg} \times L^2(\Omega)$ that is an extremal point (saddle point) of \mathcal{L} ”, or, in other words,

$$\mathcal{L}(u, p) = \tilde{J}(u) = \inf_{v \in Z_{Dg}} \tilde{J}(v) = \inf_{v \in V_{Dg}} \sup_{q \in L^2(\Omega)} \mathcal{L}(v, q) \quad (5.11)$$

The extremality conditions for \mathcal{L} are

$$d\mathcal{L}(v, 0) = \lim_{t \rightarrow 0} \frac{\mathcal{L}(u + tv, p) - \mathcal{L}(u, p)}{t} = 0 \quad \forall v \in V_{D0} \quad (5.12)$$

$$d\mathcal{L}(0, q) = \lim_{t \rightarrow 0} \frac{\mathcal{L}(u, p + tq) - \mathcal{L}(u, p)}{t} = 0 \quad \forall q \in L^2(\Omega) \quad (5.13)$$

and lead to the mixed variational formulation.

Mixed Variational Formulation of incompressible elasticity: Find $(u, p) \in V_{Dg} \times L^2(\Omega)$ such that

$$\tilde{a}(u, v) - b(v, p) = \ell(v) \quad \forall v \in V_{D0} \quad (5.14)$$

$$b(u, q) = 0 \quad \forall q \in L^2(\Omega) \quad (5.15)$$

The enforcement of incompressibility in this formulation is not built in the space for u , which is V_{Dg} and not Z_{Dg} . Instead, it appears explicitly in equation (5.15), because

$$b(u, q) = \int_{\Omega} q \operatorname{div} u \, d\Omega = 0 \quad \forall q \in L^2(\Omega) \quad \Leftrightarrow \quad \operatorname{div} u = 0 \quad \text{a.e. in } \Omega. \quad (5.16)$$

Integrating by parts the left-hand side of (5.14) one arrives at the

Differential Formulation of incompressible elasticity:

$$-\operatorname{div} \tilde{\boldsymbol{\sigma}}(u) + \nabla p = f, \quad \text{where } \tilde{\boldsymbol{\sigma}}(u) = 2\mu \boldsymbol{\epsilon}(u) \quad (5.17)$$

$$\operatorname{div} u = 0 \quad (5.18)$$

$$u = g \quad \text{on } \Gamma_D \quad (5.19)$$

$$(-p\mathbf{I} + \tilde{\boldsymbol{\sigma}}) \cdot \mathbf{n} = \mathcal{F} \quad \text{on } \Gamma_N \quad (5.20)$$

It is important to notice that the incompressibility constraint “materializes” in the equilibrium equation (5.17) as the gradient of the unknown pressure p , and at the force boundary as a normal contribution $-p\mathbf{n}$. In mechanical terms, this means that the Cauchy stress tensor of an incompressible elastic material is

$$\boldsymbol{\sigma} = -p\mathbf{I} + \tilde{\boldsymbol{\sigma}} = -p\mathbf{I} + 2\mu \boldsymbol{\epsilon}(u) \quad (5.21)$$

Exo. 5.1 Show that the extremality conditions (5.12)-(5.13) are equivalent to the mixed formulation equations (5.14)-(5.15).

Exo. 5.2 Show that, with sufficient regularity of u and p , (5.14) implies (5.17) and (5.20).

5.1.2 Constraints and mixed formulations

Most applications involve constraints: The boundary conditions themselves, as we will see, can themselves be interpreted as constraints. In Mechanics there are *kinematical constraints*, such as those imposed by some components being *rigid*, for example, or *articulated* at some places, or limited by some *obstacle*. Also, practically *any* discretization of the differential equations (5.17)-(5.18) leads to an *algebraic mixed problem*. In the *linear and finite-dimensional case*, the algebraic structure is always of the form:

$$Ax + C\lambda = f \quad (5.22)$$

$$Bx = g. \quad (5.23)$$

This means that a solution x is sought within the vectors that satisfy $Bx = g$, and that there is a *reaction* to the constraint, λ , which acts on the degrees of freedom of the system as a *force* $-C\lambda$.

It is thus important to analyze mixed problems from the algebraic point of view, dealing with *matrices* instead of with *operators* or *bilinear forms*.

Exo. 5.3 Read Chapter 3 of Mixed Finite Element Methods and Applications, by D. Boffi, F. Brezzi and M. Fortin, Springer, 2013. Then,

1. Prepare a 10-minute-long presentation of 3.1.5, Basic results. It must include a sketch of the proof of Theorem 3.1.1.

2. Prepare a 10-minute-long presentation of 3.2.1-3.2.4. It must include a sketch of the proof of Theorem 3.2.1.
3. Prepare a 10-minute-long presentation of 3.3. It must include a sketch of the proof of Theorem 3.3.1.
4. Prepare a 10-minute-long presentation of 3.4.1-3.4.2.

5.1.3 Dirichlet conditions as constraints

Up to now we have imposed the Dirichlet condition directly on the formulation space, but this condition can also be seen as a *constraint* and given a treatment similar to that given to incompressibility in the previous paragraph.

Let us illustrate this possibility in the case of a purely diffusive problem, for which the DF, PVF and PEF read

Differential Formulation:

$$-\operatorname{div}(K \nabla u) = f \text{ in } \Omega, \quad u = g \text{ on } \partial\Omega \quad (5.24)$$

Primal Variational Formulation: Find $u \in V_g = \{v \in H^1(\Omega), v = g \text{ on } \partial\Omega\}$ such that

$$a(u, v) = \int_{\Omega} K \nabla u \cdot \nabla v \, d\Omega = \int_{\Omega} f v \, d\Omega = \ell(v) \quad \forall v \in V_0 \quad (5.25)$$

Primal Extremal Formulation: Find u that extremizes J over V_g , where

$$J(v) = \frac{1}{2} a(v, v) - \ell(v) \quad (5.26)$$

To remove the Dirichlet constraint from the space, we introduce the Lagrangian

$$\mathcal{L}(v, \zeta) = J(v) - b(v - g, \zeta) \quad (5.27)$$

where

$$b(v - g, \zeta) = \int_{\partial\Omega} \zeta (v - g) \, d\Gamma = (\zeta, v - g)_{L^2(\partial\Omega)} \quad (5.28)$$

In fact, the mixed formulation that will soon be introduced is defined on a larger space than $L^2(\partial\Omega)$, denoted by $H^{-1/2}(\partial\Omega)$. The scalar product of $L^2(\partial\Omega)$ can be continuously extended as a “duality pairing” $\langle \zeta, w \rangle$ between $\zeta \in H^{-1/2}(\partial\Omega)$ and w which is in the space of traces of functions belonging to $H^1(\Omega)$, denoted by $H^{1/2}(\partial\Omega)$. Whenever ζ belongs to $L^2(\partial\Omega)$, $b(w, \zeta) = (\zeta, w)_{L^2(\partial\Omega)}$ but, in general,

$$b(w, \zeta) = \langle w, \zeta \rangle \quad (5.29)$$

Now we can write down the mixed extremal formulation of the Dirichlet problem and the mixed variational formulation that results from the corresponding extremality conditions.

Mixed Extremal Formulation: Find (u, λ) that extremizes $\mathcal{L}(\cdot, \cdot)$ over $H^1(\Omega) \times H^{-1/2}(\partial\Omega)$, i.e.,

$$\mathcal{L}(u, \lambda) = J(u) = \inf_{v \in H^1(\Omega)} \sup_{\zeta \in H^{-1/2}(\partial\Omega)} \mathcal{L}(v, \zeta) \quad (5.30)$$

Mixed Variational Formulation: Find $(u, \lambda) \in V \times Q = H^1(\Omega) \times H^{-1/2}(\partial\Omega)$ such that

$$a(u, v) - b(v, \lambda) = \ell(v) \quad \forall v \in V \quad (5.31)$$

$$b(u, \zeta) = b(g, \zeta) \quad \forall \zeta \in Q \quad (5.32)$$

and integrating by parts $a(u, v)$ we arrive at the

Mixed Differential Formulation:

$$-\operatorname{div}(K \nabla u) = f \quad \text{in } \Omega \quad (5.33)$$

$$K \frac{\partial u}{\partial n} - \lambda = 0 \quad \text{on } \partial\Omega \quad (5.34)$$

$$u = g \quad \text{on } \partial\Omega \quad (5.35)$$

which brings as new information that the Lagrange multiplier λ is in fact equal to the diffusive flux $K \partial_n u$ across $\partial\Omega$.

Exo. 5.4 *Show how to derive the mixed VF from the mixed EF.*

Exo. 5.5 *Show how to derive the mixed DF from the mixed VF.*

5.2 Abstract mixed formulation

Generalizing the previous examples, one considers the problem

Abstract Mixed Problem: Find $(u, p) \in V \times Q$ such that

$$a(u, v) - b(v, p) = \ell(v) \quad \forall v \in V \quad (5.36)$$

$$b(u, q) = g(q) \quad \forall q \in Q \quad (5.37)$$

where $a : V \times V \rightarrow \mathbb{R}$, $b : V \times Q \rightarrow \mathbb{R}$ are continuous bilinear forms, $\ell \in V'$, $g \in Q'$.

When $a(\cdot, \cdot)$ is symmetric, it is equivalent to the extremization of

$$J(v) = \frac{1}{2} a(v, v) - \ell(v) \quad (5.38)$$

over the (constrained) set

$$Z_g = \{v \in V \mid b(v, q) = g(q) \quad \forall q \in Q\} \quad (5.39)$$

and to the extremization over $V \times Q$ (i.e., unconstrained) of the Lagrangian

$$\mathcal{L}(v, q) = J(v) - b(v, q) + g(q) \quad (5.40)$$

The first logical question is whether (5.36)-(5.37) is well-posed. We consider both the cases where V and Q are infinite-dimensional (the continuous case) and finite-dimensional (the discrete case).

Theorem 5.1 *If $a(\cdot, \cdot)$ is strongly coercive on Z_0 ,*

$$a(v, v) \geq \alpha \|v\|_V^2 \quad \forall v \in Z_0 \quad (5.41)$$

with $\alpha > 0$, and if

$$\inf_{q \in Q} \sup_{v \in V} \frac{b(v, q)}{\|q\|_Q \|v\|_V} = \gamma > 0 \quad (5.42)$$

then (5.36)-(5.37) is well-posed.

The proof of this result relies on applying Thm. 1.7 to the setting defined by the product space $W = V \times Q$, the bilinear form $B : W \times W \rightarrow \mathbb{R}$ defined by

$$B((u, p), (v, q)) = a(u, v) - b(v, p) - b(u, q) \quad (5.43)$$

and the linear form $S \in W'$ defined by

$$S(v, q) = \ell(v) - g(q). \quad (5.44)$$

Exo. 5.6 *The Abstract Mixed Problem (5.36)-(5.37) is equivalent to the problem: Find $(u, p) \in W$ such that*

$$B((u, p), (v, q)) = S(v, q) \quad \forall (v, q) \in W \quad (5.45)$$

Now it only remains to prove that,

Theorem 5.2 (Brezzi) *Under hypotheses (5.41) and (5.42), the bilinear form $B(\cdot, \cdot)$ is weakly coercive on $V \times Q$.*

Proof. To simplify things, assume that (5.41) holds $\forall v \in V$ and that $a(\cdot, \cdot)$ is symmetric. Taking (u, p) arbitrary in $V \times Q$, choose $w \in V$ such that $\|w\|_V = \|p\|_Q$ and $-b(w, p) \geq \gamma\|p\|^2$. Then, taking $\eta = \alpha\gamma/N_a^2$, one gets

$$B((u, p), (u + \eta w, p)) \geq \frac{\alpha}{2} \min \left\{ 1, \frac{\gamma^2}{N_a^2} \right\} \|(u, p)\|_{V \times Q}^2$$

Besides,

$$\|(u + \eta w, p)\|_{V \times Q} \leq \left(1 + \frac{\alpha\gamma}{N_a^2} \right) \|(u, p)\|_{V \times Q}$$

so that

$$\inf_{(u,p)} \sup_{(v,q)} \frac{B((u, p), (v, q))}{\|(u, p)\| \|(v, q)\|} \geq \inf_{(u,p)} \frac{B((u, p), (u + \eta w, p))}{\|(u, p)\| \|(u + \eta w, p)\|} \geq \frac{\frac{\alpha}{2} \min \left\{ 1, \frac{\gamma^2}{N_a^2} \right\}}{1 + \frac{\alpha\gamma}{N_a^2}} > 0$$

and condition (1.22) is satisfied. Since B is symmetric, the proof is complete. As a by-product, we observe that the coercivity constant of $B(\cdot, \cdot)$ can be chosen as

$$\beta = \frac{\frac{\alpha}{2} \min \left\{ 1, \frac{\gamma^2}{N_a^2} \right\}}{1 + \frac{\alpha\gamma}{N_a^2}} \tag{5.46}$$

□

Exo. 5.7 Prove that, for all (u, p) and (v, q) in $V \times Q$,

$$B((u, p), (v, q)) \leq (N_a + 2N_b) \|(u, p)\|_{V \times Q} \|(v, q)\|_{V \times Q} \quad (5.47)$$

Exo. 5.8 Write down the abstract conditions (5.41) and (5.42) for the specific case of incompressible elastic solids or, equivalently, for incompressible viscous fluids. Sketch a proof of them in the exact case.

Hint: Consider periodic boundary conditions to simplify things. Show that proving (5.42) is equivalent to proving that, for any $q \in L_0^2(\Omega)$ (functions with zero mean), there exists $v \in H_{\text{per}}^1(\Omega)$ satisfying

$$\int_{\Omega} q \operatorname{div} v \geq c \|q\|_0^2 \quad (5.48)$$

$$\|v\|_1 \leq C \|q\|_0 \quad (5.49)$$

where c and C do not depend on q . Then define φ as the solution of $\nabla^2 \varphi = q$ and choose $v = \nabla \varphi$. Show that the two conditions above are satisfied.